

Comparative Proteomics of Excretory-Secretory Proteins Released by the Liver Fluke *Fasciola hepatica* in Sheep Host Bile and during *in Vitro* Culture *ex Host*[§]

Russell M. Morphew^{‡§}, Hazel A. Wright[‡], E. James LaCourse[¶], Debra J. Woods^{||}, and Peter M. Brophy[‡]

Livestock infection by the parasitic fluke *Fasciola hepatica* causes major economic losses worldwide. The excretory-secretory (ES) products produced by *F. hepatica* are key players in understanding the host-parasite interaction and offer targets for chemo- and immunotherapy. For the first time, subproteomics has been used to compare ES products produced by adult *F. hepatica in vivo*, within ovine host bile, with classical *ex host in vitro* ES methods. Only cathepsin L proteases from *F. hepatica* were identified in our ovine host bile preparations. Several host proteins were also identified including albumin and enolase with host trypsin inhibitor complex identified as a potential biomarker for *F. hepatica* infection. Time course *in vitro* analysis confirmed cathepsin L proteases as the major constituents of the *in vitro* ES proteome. In addition, detoxification proteins (glutathione transferase and fatty acid-binding protein), actin, and the glycolytic enzymes enolase and glyceraldehyde-3-phosphate dehydrogenase were all identified *in vitro*. Western blotting of *in vitro* and *in vivo* ES proteins showed only cathepsin L proteases were recognized by serum pooled from *F. hepatica*-infected animals. Other liver fluke proteins released during *in vitro* culture may be released into the host bile environment via natural shedding of the adult fluke tegument. These proteins may not have been detected during our *in vivo* analysis because of an increased bile turnover rate and may not be recognized by pooled liver fluke infection sera as they are only produced in adults. This study highlights the difficulties identifying authentic ES proteins *ex host*, and further confirms the potential of the cathepsin L proteases as therapy candidates. *Molecular & Cellular Proteomics* 6:963–972, 2007.

The parasitic fluke *Fasciola hepatica* infects humans and ruminant livestock worldwide. An estimated 2.4 million people are infected with *Fasciola* species, and a further 180 million are at risk (1). In addition, *F. hepatica* causes an estimated loss of \$3 billion worldwide per annum through livestock mortality, especially in sheep, and by decreased productivity via reduction of milk and meat yields in cattle (2). Presently the rate of fluke infection in parts of northwest Europe has been estimated as greater than 30%. In the absence of commercial vaccines, the benzimidazole derivative triclabendazole (TCBZ;¹ Fasinex[®]) is the drug most extensively used against *Fasciola*. Unlike other fasciolicides, TCBZ shows activity against both juvenile flukes, which are responsible for the damage to the liver of acute fasciolosis, and the mature flukes, which cause the debilitation of chronic fasciolosis. Recently resistance to TCBZ has been reported in several countries suggesting that chemotherapeutic control of this infection may soon be compromised (3–5).

The most common diagnostic field method used to detect *Fasciola* infection requires counting fluke eggs in fecal samples. However, egg counts are time-consuming and expensive and have limited ability to diagnose early acute stages of infection. New immunological diagnostic methods with commercially protected antigen identification are available in the form of a rapid on-site field test, DriDot[™] (Biotechnology Ireland) based on latex agglutination and a companion ELISA for laboratory veterinary applications. Further diagnostics-based research is required to freely validate current tests and to provide alternative molecular approaches for liver fluke populations.

In well characterized cattle models, the immune response to *F. hepatica* infection is proinflammatory lasting for about 4–6 weeks postinfection. This immune response appears to

From the [‡]Institute of Biological Sciences, University of Wales, Aberystwyth, Wales SY23 3DA, United Kingdom, ^{||}Pfizer Animal Health, Ramsgate Road, Sandwich, Kent CT13 9NJ, United Kingdom, and [¶]School of Biological Sciences, Liverpool University, Crown Street, Liverpool L69 7ZB, United Kingdom

Received, September 29, 2006, and in revised form, February 6, 2007

Published, MCP Papers in Press, February 17, 2007, DOI 10.1074/mcp.M600375-MCP200

¹ The abbreviations used are: TCBZ, triclabendazole; ES, excretory-secretory; FABP, fatty acid-binding protein; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; PMF, peptide mass fingerprint; TIC, trypsin inhibitor complex; Tpx, thioredoxin peroxidase; TTBS, Tris-buffered saline with 1% Tween 20; 2DE, two-dimensional gel electrophoresis; e, expectancy; BLAST, Basic Local Alignment Search Tool.

be switched off at around the time adults enter the bile duct. The presence of flukes modify Th1 helper T cell responses, inhibiting the induction of the protective immune system, which makes the host more susceptible to further infections. It is likely that liver fluke secretions modify host macrophage-based signaling events to switch the proinflammatory immune response “off” before the infection has been appropriately controlled.

Juvenile *F. hepatica* develop and mature within the intrahepatic bile duct and gall bladder of infected hosts, continuing to release their excretory-secretory (ES) survival products that can ultimately lead to fibrosis and calcification of host tissues. *In vitro* biochemical studies have predicted that ES products of *F. hepatica* have roles in feeding behavior, detoxification of bile components, and the evasion of the immune system. In addition, *in vitro* proteomics (6) support the release of several major protein superfamilies from liver fluke. For example, general phase II detoxification GST with proposed immune evasion roles were found to be secreted by *in vitro* cultured liver flukes (7). Multiple forms of cathepsin L proteases are also secreted *in vitro* by adult *F. hepatica* (6, 8). These proteases can cleave host hemoglobin (9) and matrix proteins (10) for nutrition and are also capable, at least *in vitro*, of cleaving host IgG (11), indicating immune evasion roles.

In vitro studies can only attempt to mimic the protein content secreted by *F. hepatica* into host bile. There is concern that the ES proteome of *F. hepatica* will be altered by parasite removal from natural host tissues and maintaining the parasite in a chemical mixture. As proteomics can identify individual proteins from a mixture of host and parasite proteins this offers the potential to validate *in vitro ex host* studies *in vivo* and provide the possibility for real time analysis. *In vivo* proteomics during parasite infection also offers the possibility of identifying new targets and validating old therapy targets, providing protein probes for sensitive and selective diagnosis and increasing our understanding of how the parasite modulates the immune system. Proteomics investigations of sheep bile have yet to be reported, and a protein-based study of human bile involved a series of chromatographic separations prior to protein identification (12, 13). Bile from domestic sheep has only been characterized previously at a gross level with respect to the major bile components (14).

Protein content in mammalian bile, including sheep, is generally within a range of 1–5 mg/ml (15). Many proteins found in the bile, such as albumin, are derived from the plasma or from the cells of the biliary system (hepatocytes and bile duct cells) and have been demonstrated in several mammalian species (16). In the gall bladder, bile is concentrated, up to 10-fold, with the content altered, often with the addition of mucus glycoproteins (16) or removal of other protein types (15). Polymeric IgA has been shown to be an important protein component of mammalian bile. However, it is not thought to be as abundant within the bile of sheep. Levels of IgG and IgA have been investigated in host bile where the host has been in-

fectured with *F. hepatica* showing considerably lower levels than that found in serum (17).

In this study proteomics approaches were used for time course analysis of protein release from *F. hepatica* during *in vitro* culture and to identify protein released from the parasite directly into the sheep bile environment. Therefore, for the first time, the ES proteome produced by *F. hepatica* into a host fluid *in vivo* was identified and compared with the *in vitro* ES proteome from *ex host* and potentially stressed parasites.

EXPERIMENTAL PROCEDURES

Bile Collection and Preparation—Gall bladders from uninfected and naturally infected sheep livers were collected immediately post-slaughter from a local abattoir. Infection status was confirmed by UK Meat Hygiene Service staff trained to identify evidence of clinical pathology associated with *Fascioliasis*. In addition, fluke infection was also confirmed immediately by the physical presence of adult *F. hepatica* within the liver and bile ducts and later by ova within extracted bile samples. Bile fluid from eight uninfected gall bladders and nine liver fluke-infected gall bladders were obtained via a sterile needle and syringe (19 gauge, 1.5 inch). Adult fluke numbers ranged from 14–65 individuals per infected sample. Care was taken to avoid removal of parasite material in infected bile samples. The individual bile samples were centrifuged at $21,000 \times g$ to clarify (removal of parasite ova and other particulate material) and then precipitated with 50% (v/v) ethanol to remove further non-protein components found in bile (16). The bile samples were further clarified via centrifugation for 15 min at $21,000 \times g$ at 4 °C. After centrifugation, the supernatant was precipitated with ice-cold 10% TCA, acetone. Bile samples were resuspended in buffer 1 (containing 6 M urea, 1.5 M thiourea, 3% (w/v) CHAPS, 66 mM DTT, 0.5% (v/v) carrier ampholytes, and protease inhibitors (CompleteMini, Roche Applied Science)).

ES Product Collection and Preparation—Live extracted *F. hepatica* collected from naturally infected sheep livers on the day of slaughter were washed for a minimum of six times in PBS, pH 7.3, at 37 °C to allow regurgitation of the parasite gut contents and to remove host material (6). ES products were collected as described previously (6) with minor alterations using 10 liver flukes per treatment replicate. Live flukes were cultured for 2, 4, 8, and 16 h at 37 °C. Care was taken to only culture intact live flukes, and all were alive when removed at the allotted time period. The ES products were clarified via centrifugation at $4000 \times g$ for 15 min at 4 °C. The ES supernatant was further centrifuged at $45,000 \times g$ for 20 min. A volume of 5 ml of ES products from each time course was precipitated with ice-cold 10% TCA, acetone. The resulting protein pellets were resuspended into buffer 2 (containing 8 M urea, 2% (w/v) CHAPS, 33 mM DTT, 0.5% (v/v) carrier ampholytes, and protease inhibitors (CompleteMini, Roche Applied Science)). At the same time as preparing ES products from live *F. hepatica*, a proportion of the extracted flukes were terminated in 1% (w/v) benzocaine in ethanol and placed in an identical medium as the live flukes. The terminated liver flukes were allowed to incubate at 37 °C for 4 h. The resulting media, containing nonspecifically released products from terminated liver flukes, were prepared exactly as the ES products from live cultured flukes as described above.

Two-dimensional Electrophoresis—A total of 300 μ l of both bile samples and ES product samples (live cultured and dead incubated), with an appropriate protein load (250 μ g for bile samples and 100 μ g for ES), were used to passively rehydrate 17-cm linear pH 3–10 IPG strips (Bio-Rad) overnight at 20 °C for separation in the first dimension. All IPG strips were focused for between 40,000 and 60,000 V-h using the Ettan IPGphor system (Amersham Biosciences). Each IPG strip was equilibrated for 15 min in 5 ml of equilibration buffer (con-

taining 50 mM Tris, pH 8.8, 6 M urea, 30% (v/v) glycerol, and 2% (w/v) SDS (18)) with the addition of DTT (Melford) at 10 mg ml⁻¹ followed by a second equilibration with iodoacetamide (Sigma) at 25 mg ml⁻¹. The IPG strips were separated in the second dimension on the Protean II system (Bio-Rad) using 12.5% polyacrylamide gels run at 40 mA for ~1 h until through a stacking gel followed by 60 mA through the resolving gel until completion. The resulting gels were Coomassie Blue-stained (PhastGel Blue R, Amersham Biosciences) and imaged via a GS-800 calibrated densitometer (Bio-Rad). Imaged 2DE gels were analyzed using Progenesis PG220 version 2006 (previously Phoretix 2D Evolution version 2005). Analysis was performed using the mode of non-spot background subtraction on average gels created from a minimum of three biological replicates. Normalized spot volumes were achieved using total spot volume multiplied by total area and were also used to determine any increase or decrease in protein abundance between comparisons (with significance set at ± 2 -fold change). Unmatched protein spots were also detected between appropriate gel comparisons. Protein spots of interest were excised and digested with trypsin (modified trypsin, sequencing grade, Roche Applied Science). Protein tryptic fragments were then eluted according to the technique of Shevchenko *et al.* (19). Samples were then resuspended in 5 μ l of 0.1% (v/v) TFA for mass spectrometry work.

Immunoblotting—*In vitro* ES and bile proteins separated via 2DE were transferred to Hybond-C nitrocellulose membrane (Amersham Biosciences) using a Trans-blot Cell (Bio-Rad) at 20 V overnight. To assess the efficiency of transfer, membranes were stained for 1 min using 0.1% Amido Black in 10% acetic acid and 25% isopropanol. Background staining was reduced using 10% acetic acid and 25% isopropanol. Following destaining, the membranes were blocked in TBS (0.1 M Tris, pH 7.5, and 0.9% (w/v) NaCl) with 1% Tween 20 (TTBS) containing 5% (w/v) skimmed milk powder for a minimum of 4 h (also removing the last traces of Amido Black staining). The blocked membranes were washed in TTBS for 10 min and then incubated with a primary antibody, either *F. hepatica*-challenged bovine serum or control/naïve bovine serum both diluted at a ratio of 1:3000 in TTBS containing 5% skimmed milk for 1 h. After three washes in TBS the membranes were incubated with a secondary antibody, anti-bovine IgG conjugated to alkaline phosphatase (Sigma) diluted to 1:30,000 in TTBS containing 5% skimmed milk for a further 1 h. Both infection serum blots and control serum blots were developed using 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate (WVR) according to the manufacturer's directions.

Mass Spectrometric Analysis—For MALDI-TOF, α -cyano-4-hydroxycinnamic acid was prepared at 10 mg ml⁻¹ in methanol. An internal standard, human angiotensin I (1296.685 Da; Sigma), was added to each sample at 1 pmol μ l⁻¹. An equal volume of sample and internal standard were mixed and added to an equal volume of α -cyano-4-hydroxycinnamic acid before 1 μ l of each was spotted onto a metal target plate and allowed to air dry. Samples were analyzed on a ToF-Spec 2E spectrometer (Waters) with delayed extraction in reflectron mode. The acquired spectra were analyzed directly using MassLynx version 3.5 (Waters) and were all externally calibrated with a mixture of peptides and lock-massed to the internal standard. Mass accuracy of the MALDI-TOF was routinely down to 10 ppm, and the resolution was consistently at 10,000 full width at half-maximum intensity. For each sample all acquired spectra were combined and processed as follows using MassLynx version 3.5: smoothing, 1 \times smooth using a Savitzky Golay method set at ± 3 channels; and background subtraction using a polynomial of order 15 and 10% below the curve to reduce background noise. To get accurate monoisotopic peak data all processed spectra were centered using the top 80% of each peak. Peak lists were generated either manually or using the ProteinLynx part of MassLynx version 3.5. For

two samples analyzed by MSMS a Q-TOF 1.5 ESI instrument (Waters) was used. Selected peptides were fragmented by CID using argon as the collision gas. Fragmentation spectra were interpreted directly using peptide sequencing (MassLynx version 3.5) following spectrum smoothing (2 \times smooths, Savitzky Golay, ± 5 channels) and processing with MaxEnt 3 deconvolution software (MassLynx version 3.5). Sequence interpretation was conducted with an intensity threshold set at 1 and a fragment ion tolerance set at 0.1 Da. Carbamidomethylation and acrylamide modifications of cysteines and oxidized methionines were taken into account.

Database Searches and Analysis—For PMF analysis, database searches were performed with the acquired monoisotopic peptide masses using MASCOT (Matrix Science). Protein identification searches allowed carbamidomethylation of cysteines as a fixed modification and oxidation of methionines, acetylation of the peptide N terminus, and conversion of glutamine to pyroglutamic acid also of the peptide N terminus as possible protein modifications. Only one maximum missed cleavage was allowed with a peptide mass tolerance of 0.2 Da. All identifications were made against the National Center for Biotechnology Information non-redundant (NCBI nr) database. Therefore, all accession numbers reported here are taken from GenBank™ (www.ncbi.nlm.nih.gov/). Searches were conducted against all metazoans or other metazoans. Signal peptide prediction analysis was performed using SignalP 3.0 Server (20) set for eukaryotes using both neural networks and hidden Markov models. Sequence alignments were performed using BioEdit version 7.0.5.3 (October 28, 2005) (21) using ClustalW multiple alignment. Peptide sequences derived from MSMS analysis were subjected to BLAST (22) to assign an identification based on sequence identity.

RESULTS

Fluke-infected and -uninfected Bile Subproteome—The bile proteomes produced from both infected and uninfected fluke individual bile samples produced reproducible 2DE arrays (Fig. 1, A and B). The matching levels using Progenesis software averaged 40% matching between replicate gels, most likely related to increases in protein trails commonly seen with serum proteins. In both infected and uninfected fluke bile samples a prominent protein cluster between the 30 and 97 kDa markers in the second dimension and between pH 5.5 and 7.5 in the first dimension was visualized along with a further protein trail spanning approximately from pH 7 to 10 at the 30-kDa marker. Image analysis of the fluke-free bile samples yielded a total of 53 protein spots present in all 2DE arrays. The fluke-infected bile array showed a total of 60 protein spots consistently visualized on all 2DE arrays. A cluster of 15 protein spots located at the 30-kDa marker and spreading from approximately pH 5 to 6 was visible in all the fluke-infected bile samples; these protein spots were absent in the fluke-free bile samples (15 of 23 spots in total from pH 3 to 10 shown in Fig. 1C).

***In Vitro* ES Product Time Course Proteomes**—The ES products released during *F. hepatica in vitro* culture were separated by 2DE after 2, 4, 8, and 16 h postculture (Fig. 2). The *in vitro* derived 2DE profiles were compared with the *in vivo* bile liver fluke ES subproteome profile (Fig. 1) to validate the applicability of *in vitro* culture methods currently used by parasitic worm research communities. For up to 8 h *ex host*, the *in vitro* culture 2DE profile was similar to the *in vivo* profile

FIG. 1. **Representative 2DE protein arrays of infected and uninfected host bile proteomes for *in vivo* analysis.** Proteins were separated across a linear pH range of 3–10 using IEF in the first dimension and 12.5% SDS-PAGE in the second dimension and Coomassie Blue-stained. *A*, *F. hepatica*-infected sheep bile protein array. *B*, *Fasciola*-free/uninfected sheep bile proteome. *C*, Progenesis analysis identifying major changes between both proteomes, here showing 23 outlined spots unique to the infected bile proteome. Of these 23 unique proteins, 10 identifications corresponded to proteins originating from the parasite, and one identification corresponded to protein originating from the host (Table I). In *A*, *B*, and *C* numbered and circled protein spots correspond to putative identifications located in Table I.





FIG. 2. **Representative 2DE protein arrays of *F. hepatica* ES products from *in vitro* culture.** Proteins were separated across a linear pH range of 3–10 using IEF in the first dimension and 12.5% SDS-PAGE in the second dimension and Coomassie Blue-stained. ES products are from 2 h in culture (*A*), 4 h in culture (*B*), 8 h in culture (*C*), and 16 h in culture (*D*). Numbered and circled spots correspond to putative identifications located in Table I.

of infected bile samples (Fig. 2), although a limited proteome was produced of which the majority of the protein spots visualized migrated to the 30-kDa marker between pH 5 and 6. Overall complexity of the *in vitro* ES profiles increased after 8 h *ex host* with the most noticeable increases present between 45 and 66 kDa. In particular two protein spots, spot 16 estimated at 49 kDa and spot 17 estimated at 46 kDa (Fig. 2), showed significant increases after 16 h in culture (Fig. 3; spot

16 showed a 2.4-fold increase, and spot 17 showed a 3.6-fold increase). In addition, two protein spots (Fig. 2, two spots both identified as spot 13) present between 14.4 and 20 kDa, estimated at 14.8 kDa using Progenesis, also increased over time (Fig. 3; after 8 h of culture a 3.8-fold increase for one and a 2.8-fold increase for the other). In relation to previous ES studies on *F. hepatica* the cluster of protein spots present at the 30-kDa marker are in agreement with that found by Jef-




FIG. 3. Montage views from Progenesis PG220 version 2006 analysis of *in vitro* culture ES products visualizing increases in abundance of three *F. hepatica* proteins, actin, enolase, and FABP (present in two locations). Actin (spot 17) and enolase (spot 16) both showed significant increases, 2.4- and 3.6-fold increases, respectively, in abundance within the *in vitro* culture array. FABP, represented as two spots on the 2DE array (spots labeled 13), also showed a significant increase in abundance, although significantly more is seen after only 8 h in culture (3.8- and 2.8-fold increases after 8 h). All spot numbers relate to putative identifications located in Table I. Spot numbers also correspond directly to those in Fig. 2.

feries *et al.* (6). However, the remaining proteins seen by Jefferies *et al.* (6) only begin to appear after time, especially in the 16-h profile. All of the ES profiles analyzed were reproducible (80–94% matching between *in vitro* ES product replicates) as only viable flukes were taken to produce accurate 2DE *in vitro* assays. Analysis of the dead fluke cultured ES proteome (Fig. 4) showed an increase in total overall protein spots visualized and yielded a total of 107. Matching between live ES products and the dead ES proteome was relatively poor when compared with the live ES products (13.1% matching *versus* 2-h ES proteome and 22.4% matching *versus* 16-h proteome) and was related to the increase in protein spots detected. The large cluster of proteins observed between pH 5 and 6 and migrating to the 30-kDa marker in live ES proteomes was vastly reduced in the dead ES proteome with large increases in protein abundance seen all over the remainder of the array.

Identification of Protein Spots from *in Vivo* and *in Vitro* Approaches—A total of 90 protein spots from all assays were cut from gels for identification via PMF. This included spots from fluke-infected (20 spots) and fluke-free bile proteomes

(10 spots as marker proteins) and *in vitro* culture proteomes (25 spots from the 16-h profile and 35 spots from the dead ES proteome). For identification through MASCOT several factors were exploited to assign a significant identification, *i.e.* expectancy (*e*) value, percent coverage, M_r , *pI*, and the error expressed in ppm. MASCOT also has the added advantage of being able to assign statistical significance of which a minimum of $p = 5\%$ was taken (wherever possible significance was increased to $p = 1\%$ or $p = 0.1\%$). Positive identifications were made on 67 occasions (Table I) with 15 from infected bile proteomes (Fig. 1A), six from uninfected bile proteomes (Fig. 1B), 21 from *in vitro* culture proteomes (Fig. 2), and a further 25 from the dead ES proteome (Fig. 4).

From both bile proteomes, six proteins were identified as host proteins, potential bile-based biomarkers of fluke infection. Three of the identifications were based on significant hits on *Bos taurus* due to the lack of or limited sequences for *Ovis aries*. Two of these identifications based on *B. taurus*, transferrin and enolase, were subjected to sequence alignment with available *O. aries* sequence to further support identification. Further verification of enolase as a host protein and not

parasite enolase was required. To confirm enolase as a host protein the short sequence of *O. aries* (GenBank accession number AAF60279), the *B. taurus* (GenBank accession number NP_776474) sequence hit from PMF, the top enolase

BLAST (22) hit sequence from *B. taurus* (GenBank accession number AAI02989), and the top *F. hepatica* (GenBank accession number AAF60279) BLAST hit were subjected to sequence alignment. Matching peptides (most importantly the peptide at *m/z* 1556.8; sequence VVIGMDVAASEFYR) from PMF analysis corresponded to conserved sequences from both *B. taurus* sequences and that of the *O. aries* sequence but not that of *F. hepatica* confirming identification of enolase as a host derived protein.² The final identification based on *B. taurus* sequences was that of regucalcin with no sequence available for *O. aries*; identity was assigned by the e value of a protein highly conserved in mammals (23).

Of the 15 identified proteins found in *Fasciola*-infected bile 10 were identified as *Fasciola* cathepsin L proteases. However, only two of these identifications could be significantly assigned to a specific entry in the public database, confirming problems in proteomics identification by PMF (6). The two significant identifications corresponded to a cathepsin L-like protease (GenBank accession number CAA80446) and secreted cathepsin L2 (GenBank accession number AAC47721) (Fig. 1, spots 1 and 2, respectively). These two could be distinctly separated from the remaining cathepsin hits due to unique peptides appearing in the PMFs, for CAA80446 a peptide of *M_r* 2449.07⁺ (monoisotopic) corresponding to the peptide DQGQCGSCWAFSTTGAVEGQFR and for AAC47721 a peptide of *M_r* 2448.05⁺ (monoisotopic) corresponding to NQGQCGSCWAFSTTGAVEGQFR. Due to the high conserva-




FIG. 4. Representative 2DE protein array of *F. hepatica* proteins nonspecifically released from *in vitro* incubation of dead adults. Proteins were separated across a linear pH range of 3–10 using IEF in the first dimension and 12.5% SDS-PAGE in the second dimension and Coomassie Blue-stained. Numbered and circled spots correspond to putative identifications located in Table I.

² Data not shown.

TABLE I

Putative identifications of proteins revealed in both *in vivo* bile and *in vitro* ES proteomes utilizing PMF

MASCOT e values significant at *p* = 5% (*), *p* = 1% (**), and *p* = 0.1% (***) are shown. SP, signal peptide predicted using SignalP. WB, showed immunogenic properties during Western blotting. Av, average. All spot identification numbers (1–18 and A and B) correlate across figures, for example, spot 1 in Fig. 1 is the same cathepsin L identified as spot 1 in Figs. 2 and 4.

Spot N°	MASCOT (e)	Sequence Coverage (%)	Matched Peptides (Total)	Progenesis Calculated <i>M_r</i> / <i>pI</i>	Theoretical <i>M_r</i> / <i>pI</i>	GenBank Accession N°	Species	Description	Where Identified	Secreted Protein
1	0.003**	39	12(30)	24118 / 4.87	37534 / 5.43	CAA80446	<i>F. hepatica</i>	Cathepsin L-like proteinase	<i>In vivo In vitro</i>	SP, WB
2	0.011**	43	16(43)	24066 / 5.00	37466 / 5.54	AAC47721	<i>F. hepatica</i>	Secreted Cathepsin L 2	<i>In vivo In vitro</i>	SP, WB
3	0.5	25	9(39)	24497 / 5.16	37524 / 5.55	AAF76330	<i>F. hepatica</i>	Cathepsin L	<i>In vivo In vitro</i>	SP, WB
4	Av. 0.88	Av. 25	Av. 9(26)	23741-23825 / 5.55-5.93	35611 / 5.79	AAK38169	<i>F. hepatica</i>	Cathepsin L-like	<i>In vivo In vitro</i>	WB
5	Av. 7.6	Av. 17	Av. 5(20)	23588 / 6.50	24600 / 5.01	AAF44678	<i>F. gigantica</i>	Cathepsin L	<i>In vivo In vitro</i>	WB
6	Av. 0.93	Av. 43	Av. 8(34)	24485 / 6.48	35567 / 5.79	AAM11647	<i>F. hepatica</i>	Cathepsin L	<i>In vitro</i>	WB
7	0.031*	65	9(43)	17854 / 6.50	15725 / 6.16	P09670	<i>O. aries</i>	Superoxide dismutase [Cu-Zn]	<i>In vivo</i>	
8	1.4E-05***	64	13(48)	26857 / 7.01	29079 / 6.4	P00922	<i>O. aries</i>	Carbonic anhydrase II (Carbonate dehydratase II) (CA-II)	<i>In vivo</i>	
9	0.00043***	41	15(46)	31470 / 5.97	33308 / 5.54	NP_776382	<i>B. taurus</i>	Regucalcin (Senescence marker protein-30)	<i>In vivo</i>	
10	0.00081***	42	13(36)	48014 / 6.71	47277 / 6.44	NP_776474	<i>B. taurus</i>	Enolase 1	<i>In vivo</i>	
11	1.8E-11***	49	27(57)	74794 / 6.19	69188 / 5.80	P14639	<i>O. aries</i>	Serum albumin precursor	<i>In vivo In vitro</i> ¹	
12	0.016*	24	15(38)	80914 / 6.86	77753 / 6.75	NP_803450	<i>B. taurus</i>	Transferrin	<i>In vivo</i>	
13	8.9E-05***	69	12(34)	17039 / 6.17	14805 / 5.95	Q7M4G1	<i>F. hepatica</i>	Fatty Acid-Binding Protein Type II	<i>In vitro</i>	
14	0.03*	46	15(50)	23812 / 7.15	25373 / 5.89	AAB28746	<i>F. hepatica</i>	Glutathione s-transferase 26 KD 51 (GST51) (FH51) (GST Class-Alpha)	<i>In vitro</i>	
15	0.0044**	49	13(40)	34758 / 8.21	23687 / 7.04	AAG23287	<i>F. hepatica</i>	Glyceraldehyde Phosphate Dehydrogenase	<i>In vitro</i>	
16	8.9E-07***	48	22(50)	49231 / 6.95	46707 / 6.56	A53665	<i>F. hepatica</i>	Phosphopyruvate Hydratase (Enolase)	<i>In vitro</i>	
17	0.0047**	28	10(65)	46335 / 5.95	41157 / 5.56	AAAB2603	<i>D. dendriticum</i>	Actin	<i>In vitro</i>	
18	0.0013**	57	9(24)	17846 / 9.45	14671 / 9.02	CAB65015	<i>F. hepatica</i>	Fatty Acid Binding Protein Type III	<i>In vitro</i> ¹	
A, B	4e-06 ^a	13	3(3) ^b WVYSAAH LQGVSWGYGCAAGK GRSNGGINVAEGNEKFIASAK	25429-26101 / 7.65-9.29	23305 / 8.69	1EB2_A	<i>B. taurus</i>	Chain A, Trypsin Inhibitor Complex (Bpo)	<i>In vivo</i>	WB ^c

^a BLAST score derived from searches using MSMS data.

^b Peptides sequence using MSMS; underlined regions indicate matches to the top scoring sequence.

^c TIC was identified as a host protein; although it was recognized by Western blotting, it is not derived from *F. hepatica*.

¹ Serum albumin and FABP type III were identified *in vitro* in dead ES products only.




FIG. 5. **Western blot analysis of ES products from *in vivo* analysis.** Transferred proteins were probed with *Fasciola*-challenged and *Fasciola*-naïve bovine sera to assess the antigenicity of the protein present in bile to identify true secreted proteins. *A*, *Fasciola*-infected bile probed with *Fasciola*-challenged serum. *B*, *Fasciola*-infected bile probed with *Fasciola*-naïve serum, demonstrating the reactivity of the secondary antibody (anti-bovine IgG) to many of the host proteins within the bile proteome. Areas enclosed by a box indicate proteins displaying antigenic properties and were identified as *F. hepatica* cathepsin L proteases. Circled spots outline a major contributor to the Western blot profile later identified as trypsin inhibitor complex (Table I, spots A and B).

tion between cathepsin L sequences all the other cathepsin hits could not be identified to a specific entry, although all were highest scoring hits. Interestingly one cathepsin L identification (Fig. 1, spot 5) top scored to a cathepsin L from the closely related tropical liver fluke, *Fasciola gigantica* (GenBank accession number AAF44678).

As *in vivo*, cathepsin L proteases could be identified *in vitro* but in this instance in 15 locations. Only two of these 15 cathepsin L proteases could be significantly identified to a specific entry in the public databases, appearing only once within the ES profile both *in vivo* and *in vitro* (GenBank accession numbers CAA80446 and AAC47721; Fig. 2, spots 1 and 2, respectively). In both *in vivo* and *in vitro* preparations the cathepsin L, GenBank accession number AAF76330 (Figs. 1 and 2, spot 3), was assigned to the same location but was not significantly confirmed. Two further cathepsin L hits assigned using *in vitro* methods were again from *F. gigantica* (GenBank accession number AAF44678; Fig. 2, spots labeled 5). The remaining proteins identified from *in vitro* cultures, with the exception of one, were assigned to proteins from *F. hepatica* sequences. The one protein assigned to an entry not of *F. hepatica* was that of the structural protein actin from the pseudophyllidean cestode *Diphyllobothrium dendriticum*. There is high conservation among actin sequences between closely related organisms, and with a highly significant score at $p = 0.1\%$, this hit was assumed to be a form of actin. Analysis of the dead ES proteins again identified those seen after 16-h culture including actin (based on *D. dendriticum*) and enolase although in greater numbers (*i.e.* actin, four locations; enolase, three locations; and fatty acid-binding protein (FABP) type II, four locations). In addition to this, a FABP was identified at high pH, estimated at pI 9.45, and was only

identified in dead ES. The relative contribution of the cathepsin L proteases to the dead ES subproteome was vastly reduced. Interestingly serum albumin from the host was identified in dead ES but not in live cultures.

Immunoblotting of *in Vitro* and *in Vivo* ES Proteomes of *F. hepatica*—Western blotting with *Fasciola*-challenged and naïve bovine sera as the probes was used to compare the antigenicity pattern of the *in vivo* (Fig. 5) and *in vitro*³ ES proteomes of *F. hepatica*. Despite some cross-reactivity with the secondary antibody, anti-bovine IgG, antigenic proteins from *in vivo* analysis (most likely from *F. hepatica* secretions) could be distinguished. In this manner 27 protein spots were identified as antigenic. Of these 27, 11 were associated with trains of protein that also cross-reacted in the control, leaving a total of 16 that were truly antigenic. Identification of 15 of these antigens corresponded to the *F. hepatica* cathepsin L proteases secreted into the bile.

Two protein spots contributed significantly to the Western blots from *F. hepatica*-infected and, to a lesser extent, uninfected bile proteomes. After failure to identify them using PMF analysis, both spots were subjected to peptide sequencing using MSMS. Following BLAST analysis both were identified as bovine trypsin inhibitor complex (TIC; GenBank accession number 1EB2_A) shown in Table I and Fig. 1, spots A and B. Notably TIC protein spot A was increased in *F. hepatica*-infected bile proteomes by a factor of 3.98, and spot B was absent in control bile. Western blotting of *in vitro* cultured ES products³ allowed this profile to be further confirmed for *in vivo* secretion and exposure to host immune system. Of all of

³ Data not shown. Western blotting of *in vitro* 16-h ES products revealed only 15 cathepsin L proteases as immunogenic.

the protein spots in the 16-h profile of ES products, 15 were shown to produce antigenic responses. Again following identification, all 15 were shown to be cathepsin L proteases produced by fluke *in vitro*. No other proteins displayed antigenic properties. Immunoblotting of the dead ES proteome² again revealed the cathepsin L proteases as immunogenic when probed with *Fasciola*-challenged serum. Actin and enolase were also recognized but by naïve control serum.

Analysis of all the parasite sequences derived from PMF identifications for signal peptide cleavage sites revealed only a small fraction of the *in vivo* and *in vitro* profiles to be secretory. This corresponded to three GenBank protein entries (accession numbers CAA80446, AAC47721, and AAF76330) all of which are *F. hepatica* cathepsin L proteases. No other protein sequences analyzed showed potential cleavage sites for signal peptides.

DISCUSSION

A rapid sample preparation strategy for undertaking reproducible 2DE protein arrays from sheep bile from *F. hepatica*-infected individuals has been optimized. In addition, without completed and verified genome support, proteins were identified by PMF from the sheep host, *O. aries*, and for the first time *in vivo* its parasite, the liver fluke *F. hepatica*. This is the first experimental *in vivo* identification of ES products from an adult parasitic worm living in a mammalian host. Our *in vivo* host bile analysis suggests that the major *F. hepatica* proteins present outside the parasite in host tissues were a variety of proteases, mainly the cathepsin L proteases. This supports previous *in vitro* studies, suggesting that these proteins play a major role in *F. hepatica* survival in the host gall bladder and bile duct. *In vivo* proteomics supports the suggested role of cathepsins in providing nutrition with the secreted cathepsin L2 (Figs. 1 and 2, spot 2) cleaving fibrinogen to form blood clots and prevent excessive bleeding at feeding sites (24) and the cathepsin L-led disruption of the immune system via cleavage of host immunoglobulin (25) commonly circulating within the bile (17). These noted differences in enzymatic properties seen in cathepsin L proteases may well be related to changes of amino acid composition in and around the active site where Irving *et al.* (26) have detailed many sites that are subjected to positive selection pressures.

Cathepsin L proteases were the only *F. hepatica* proteins identified in both bile and *in vitro* culture. In addition, our *in vitro* studies confirm that cathepsin L proteases are the major adult ES proteins to at least 16 h postculture. GST (Type 51), FABP (type II), and enolase were also continually released from *F. hepatica* for at least 16 h during *in vitro* ES culture. These proteins have all been detected previously *in vitro* ES products from *F. hepatica* (6, 27, 28) but were not observed our *in vivo* assays. In addition, *F. hepatica* GST (Type 51), FABP (type II), and enolase have no obvious signal sequence and are not recognized by pooled infection sera. Although GSTs were not identified *in vivo*, the levels of glutathione in

the bile (1–4 mM in mammals with 90% in the reduced form (29–31)) would allow for GST enzymatic activity (K_m of GSTs, ~0.4 mM) suggesting that the stress-responsive detoxification protein GST could, in theory, function on the surface of the fluke. The known vaccine potential of *F. hepatica* GST may relate to an anti-inflammatory GST being secreted by newly excysted juveniles that are first exposed to the immune system. Enolase has been found on the surface of many other pathogenic species, *i.e.* bacteria, fungi, and protozoa (32) with immune modulating properties (33), and has been predicted on the surface of the nematode worm *Onchocerca volvulus* (34) and found in *in vitro* culture studies in other nematodes, *Ascaris suum* (35) and *Haemonchus contortus* (36).

Several proteins showed significant increases in the *in vitro* system at 16 h postculture, including FABP, actin, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH). We believe these proteins are *in vitro* artifacts, especially given the lack of *in vivo* validation and absence of immune recognition. We suggest that FABP probably functions in intraparasite lipid transport in adult *F. hepatica* despite being found in ES preparations in previous *in vitro* studies on *F. hepatica* (6) and *Schistosoma mansoni* (37). Bile contains high levels of fatty acids that allow the parasite to avoid synthesis and simply take up fatty acids from the host (38). FABP has been located to lipid droplets below the subtegumental region of adult male and female *Schistosoma japonicum* flatworms but was not present at the surface, suggesting a role in fatty acid transport not uptake *per se* (38). GAPDH is a major surface antigen in both larval stages and the adult of the blood fluke *S. mansoni* (39, 40). Therefore, GAPDH may also be present at the surface of the related flatworm *F. hepatica* at least in the adult life cycle stage.

Several host proteins were also identified despite an incomplete genome, taking advantage of cross-species identification (41). Some of these host proteins have been found previously in other bile-based studies (13) and have well characterized functions, such as carbonic anhydrase II (balance and regulation of acid-base levels; Ref. 42), transferrin (iron transport protein) delivered to the intestine via the bile duct system (43), and albumin (a secretory product of the liver used to transport large organic anions; Ref. 44). The glycolytic enzyme enolase was confirmed as a host protein *in vivo* as others have suggested it is a *Fasciola* ES protein (28). Sheep TIC identified in host bile has been shown to be highly immunogenic (45). TIC inactivates excessive trypsin, and its increase in liver fluke-infected bile may be part of a defense to directly counteract parasite survival roles of *F. hepatica* proteases. An immunogenic TIC may function as a host biomarker for liver fluke infection via antibody-based assays in bulk milk, blood, or feces. This would support current blood-based tests, DriDot, and its companion ELISA that rely on an *F. hepatica* protein to enter the blood circulation via feeding on damaged host tissues. Others are exploring SELDI tech-

nology for parasite (and host) blood-borne biomarkers.⁴

Thus, why are GST, FABP, enolase, actin, and GAPDH released by *F. hepatica in vitro* and are not detected *in vivo*? Adult *F. hepatica* are thought to continually shed their teguments to evade the host immune system (46). As GST, FABP, enolase, actin, and GAPDH are located at the surface or just below the surface of the fluke, they may be released into the surrounding culture medium as the tegument is sloughed. As *in vitro* time progresses in *ex host* stress response-inducing environments, adult flukes may shed their teguments at a greater rate. Tegumental turnover at an elevated rate facilitates the release of GST, FABP, enolase, actin, and GAPDH. In a schistosome flatworm proteomics study GST, FABP, enolase, GAPDH, and actin also were identified in soluble extracts of isolated *S. mansoni* teguments (47). The ES proteome from dead cultured *F. hepatica* fluke further confirms that the release of these proteins *in vitro* is not a specific biological secretion but a physical degradation process. Although found in previous ES studies (6, 48, 49) the predicted helminth antioxidant, immunosuppressant, and vaccine candidate thioredoxin peroxidase (Tpx) was not identified in this study using *in vivo* or *in vitro* techniques. In the flatworm *S. mansoni*, Tpx was located in soluble tegumental preparations and located beneath the surface (47). This indicates that cultured adult liver flukes may need to shed their tegument continuously before Tpx will be seen in ES products or that it results from degradation of cultured flukes. Tpx may be on the surface or secreted by juvenile stages of *F. hepatica* that are more exposed to the immune response, hence its protection properties during vaccine trials (50). As our study suggests, both FABP and GST are not proactively secreted *in vivo* by adult *F. hepatica* but released via shedding, questioning their inclusion in vaccine formulations or as anti-inflammatory immune modulators as they only appear to be exposed intermittently during shedding.

The ability of proteomics to discriminate between parasite and host in biological fluids, such as in the bile via percutaneous cholecystocentesis (16, 51), will increase understanding of how helminth worms regulate the immune system to establish chronic infections. Other host fluids where *F. hepatica* reside may also be suitable to this type of proteomics analysis. Newly excysted juveniles emerge from their metacercarial cysts in the duodenum where early ES products are released, and migrating juveniles pass through the peritoneal cavity on route to the liver and will again release ES products into the surrounding host fluid. However, in both of these instances the relative abundance of newly excysted juveniles and juvenile ES products in relation to host protein may severely hamper proteomics analysis of these host fluids. Nonetheless the analysis of *in vivo* ES products from intestinal helminth worms may be possible by analyzing intestinal fluid or washes.

⁴ T. W. Spithill, personnel communication.

In vivo proteomics technology has identified for the first time proteins released from the liver fluke *F. hepatica*, a prerequisite to understanding the host-adult parasite interface of this important global parasite. This approach also supports the academic portfolio of the cathepsin L proteases as liver fluke therapeutic candidates and suggests the need to be careful extrapolating findings from *ex host in vitro* studies. *In vivo* proteomics presents an opportunity to validate *in vitro* findings, such as cathepsin L cleavage of host IgG molecules, and although this particular assay was not completed in the present study the potential applications of our technique is demonstrated. Our present study only provides a snapshot of the interaction between host and parasite; a live bile extraction, via a cannula (16, 51), offers opportunities for time course analysis of an experimental infection. The identification of *F. hepatica* proteins in the host from relatively low infection levels (60 flukes) may be translated to biomarker analysis to support current *F. hepatica* infection diagnosis.

Acknowledgments—We thank Dr. Jim Jefferies for technical assistance; Jim Heald and Charly Morgan for MS skills; the staff at Dunbia (Llanybydder, Wales, UK), in particular Gerwyn Probert, for assistance; and Professor Grace Mulcahy (Dublin, Ireland) for providing the bovine *Fasciola*-challenged and naïve sera.

* This work was supported by Biotechnology and Biological Sciences Research Council (BBSRC) Grant C503638/2, European Union (DELIVER: Grant FOOD-CT-200X-023025), and Pfizer Animal Health Co-operative Awards in Science and Engineering (CASE) (BBSRC Committee Ph.D. Grant BBS/S/P/2003/10460). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

§ The on-line version of this article (available at <http://www.mcponline.org>) contains supplemental material.

§ To whom correspondence should be addressed. Tel.: 44-1970-622337; Fax: 44-1970-622350; E-mail: rmm97@aber.ac.uk.

REFERENCES

1. Anonymous (1995) *Control of Foodborne Trematode Infections*, World Health Organization, Geneva
2. Boray, J. C. (1994) Chemotherapy of infections with fasciolidae, in *Immunology, Pathobiology and Control of Fasciolosis* (Boray, J. C., ed) pp. 83–97, MSD AGVET, Rahway, NJ
3. Moll, L., Gaasenbeek, C. P. H., Vellema, P., and Borgsteede, F. H. M. (2000) Resistance of *Fasciola hepatica* against triclabendazole in cattle and sheep in The Netherlands. *Vet. Parasitol.* **91**, 153–158
4. Mitchell, G. B. B., Maris, L., and Bonniwell, M. A. (1998) Triclabendazole-resistant liver fluke in Scottish sheep. *Vet. Rec.* **143**, 399–399
5. Thomas, I., Coles, G. C., and Duffus, K. (2000) Triclabendazole-resistant *Fasciola hepatica* in southwest Wales. *Vet. Rec.* **146**, 200–200
6. Jefferies, J. R., Campbell, A. M., Van Rossum, A. J., Barrett, J., and Brophy, P. M. (2001) Proteomic analysis of *Fasciola hepatica* excretory-secretory products. *Proteomics* **1**, 1128–1132
7. Brophy, P. M., and Barrett, J. (1990) Glutathione transferase in helminths. *Parasitology* **100**, 345–349
8. Wijffels, G. L., Panaccio, M., Salvatore, L., Wilson, L., Walker, I. D., and Spithill, T. W. (1994) The secreted cathepsin L-like proteinases of the trematode, *Fasciola hepatica*, contain 3-hydroxyproline residues. *Biochem. J.* **299**, 781–790
9. Dalton, J. P., Neill, S. O., Stack, C., Collins, P., Walshe, A., Sekiya, M., Doyle, S., Mulcahy, G., Hoyle, D., Khaznadji, E., Moire, N., Brennan, G.,

- Mousley, A., Kreshchenko, N., Maule, A. G., and Donnelly, S. M. (2003) *Fasciola hepatica* cathepsin L-like proteases: biology, function, and potential in the development of first generation liver fluke vaccines. *Int. J. Parasitol.* **33**, 1173–1181
10. Berasain, P., Goni, F., McGonigle, S., Dowd, A., Dalton, J. P., Frangione, B., and Carmona, C. (1997) Proteinases secreted by *Fasciola hepatica* degrade extracellular matrix and basement membrane components. *J. Parasitol.* **83**, 1–5
11. Smith, A. M., Dowd, A. J., Heffernan, M., Robertson, C. D., and Dalton, J. P. (1993) *Fasciola hepatica*: a secreted cathepsin L-like proteinase cleaves host immunoglobulin. *Int. J. Parasitol.* **23**, 977–983
12. Stark, M., Jornvall, H., and Johansson, J. (1999) Isolation and characterization of hydrophobic polypeptides in human bile. *Eur. J. Biochem.* **266**, 209–214
13. Kristiansen, T. Z., Bunkenborg, J., Gronborg, M., Molina, H., Thuluvath, P. J., Argani, P., Goggins, M. G., Maitra, A., and Pandey, A. (2004) A proteomic analysis of human bile. *Mol. Cell. Proteomics* **3**, 715–728
14. Kaur, B., and Ahuja, S. P. (1993) Characterization of biliary proteins and composition of bile from sheep and goats. *Zentbl. Vetmed. Reihe A* **40**, 598–604
15. La Russo, N. F. (1984) Proteins in bile—how they get there and what they do. *Am. J. Physiol.* **247**, G199–G205
16. Coleman, R. (1987) Biochemistry of bile secretion. *Biochem. J.* **244**, 249–261
17. Ferre, I., OrtegaMora, L. M., and RojoVazquez, F. A. (1997) Serum and bile antibody responses (IgG and IgA) during subclinical *Fasciola hepatica* infection in sheep. *Vet. Parasitol.* **68**, 261–267
18. Gorg, A., Postel, W., Weser, J., Gunther, S., Strahler, J. R., Hanash, S. M., and Somerlot, L. (1987) Elimination of point streaking on silver stained two-dimensional gels by addition of iodoacetamide to the equilibration buffer. *Electrophoresis* **8**, 122–124
19. Shevchenko, A., Wilm, M., Vorm, O., and Mann, M. (1996) Mass spectrometric sequencing of proteins from silver stained polyacrylamide gels. *Anal. Chem.* **68**, 850–858
20. Bendtsen, J. D., Nielsen, H., von Heijne, G., and Brunak, S. (2004) Improved prediction of signal peptides: SignalP 3.0. *J. Mol. Biol.* **340**, 783–795
21. Hall, T. A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* **41**, 95–98
22. Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410
23. Misawa, H., and Yamaguchi, M. (2000) The gene of Ca²⁺-binding protein regucalcin is highly conserved in vertebrate species. *Int. J. Mol. Med.* **6**, 191–196
24. Dalton, J. P., McGonigle, S., and Dowd, A. J. (1994) Cathepsin L2 secreted by the parasitic trematode *Fasciola hepatica* cleaves fibrinogen in a novel manner that produces a fibrin clot. *J. Cell. Biochem.* **18D**, 166–166
25. Berasain, P., Carmona, C., Frangione, B., Dalton, J. P., and Goni, F. (2000) *Fasciola hepatica*: parasite-secreted proteinases degrade all human IgG subclasses: determination of the specific cleavage sites and identification of the immunoglobulin fragments produced. *Exp. Parasitol.* **94**, 99–110
26. Irving, J. A., Spithill, T. W., Pike, R. N., Whisstock, J. C., and Smooker, P. M. (2003) The evolution of enzyme specificity in *Fasciola* spp. *J. Mol. Evol.* **57**, 1–15
27. Cervi, L., Rossi, G., and Masih, D. T. (1999) Potential role for excretory-secretory forms of glutathione-S-transferase (GST) in *Fasciola hepatica*. *Parasitology* **119**, 627–633
28. Bernal, D., de la Rubia, J. E., Carrasco-Abad, A. M., Toledo, R., Mas-Coma, S., and Marcilla, A. (2004) Identification of enolase as a plasminogen-binding protein in excretory-secretory products of *Fasciola hepatica*. *FEBS Lett.* **563**, 203–206
29. Ballatori, N., and Clarkson, T. W. (1982) Developmental changes in the biliary excretion of methylmercury and glutathione. *Science* **216**, 61–63
30. Ballatori, N., and Clarkson, T. W. (1983) Biliary transport of glutathione and methylmercury. *Am. J. Physiol.* **244**, G435–G441
31. Eberle, D., Clarke, R., and Kaplowitz, N. (1981) Rapid oxidation *in vitro* of endogenous and exogenous glutathione in bile of rats. *J. Biol. Chem.* **256**, 2115–2117
32. Pancholi, V. (2001) Multifunctional α -enolase: its role in diseases. *Cell. Mol. Life Sci.* **58**, 902–920
33. Ferreira, P., Bras, A., Tavares, D., Vilanova, M., Ribeiro, A., Videira, A., and AralaChaves, M. (1997) Purification, and biochemical and biological characterization of an immunosuppressive and lymphocyte mitogenic protein secreted by *Streptococcus sobrinus*. *Int. Immunol.* **9**, 1735–1743
34. Jolodar, A., Fischer, P., Bergmann, S., Buttner, D. W., Hammerschmidt, S., and Brattig, N. W. (2003) Molecular cloning of an α -enolase from the human filarial parasite *Onchocerca volvulus* that binds human plasminogen. *Biochim. Biophys. Acta* **1627**, 111–120
35. Andrade, M. A., Siles-Lucas, M., Lopez-Aban, J., Carranza, C., Perez-Arellano, J. L., and Muro, A. (2005) Antigens from *Ascaris suum* trigger *in vitro* macrophage NO production. *Parasite Immunol.* **27**, 235–242
36. Yatsuda, A. P., Krijgsveld, J., Cornelissen, A., Heck, A. J. R., and de Vries, E. (2003) Comprehensive analysis of the secreted proteins of the parasite *Haemonchus contortus* reveals extensive sequence variation and differential immune recognition. *J. Biol. Chem.* **278**, 16941–16951
37. Thaumaturgo, N., Vilar, M. M., Diogo, C. M., Edelenyi, R., and Tendler, M. (2001) Preliminary analysis of Sm14 in distinct fractions of *Schistosoma mansoni* adult worm extract. *Mem. Inst. Oswaldo Cruz* **96**, 79–83
38. Timanova-Atanasova, A., Jordanova, R., Radoslavov, G., Deevska, G., Bankov, I., and Barrett, J. (2004) A native 13-kDa fatty acid binding protein from the liver fluke *Fasciola hepatica*. *Biochim. Biophys. Acta* **1674**, 200–204
39. Trudgett, A., McNair, A. T., Hoey, E. M., Keegan, P. S., Dalton, J. P., Rima, B. K., Miller, A., and Ramasamy, P. (2000) The major tegumental antigen of *Fasciola hepatica* contains repeated elements. *Parasitology* **121**, 185–191
40. Goudot-Crozal, V., Caillol, D., Djabali, M., and Dessein, A. J. (1989) The major parasite surface antigen associated with human resistance to Schistosomiasis is a 37-kD glyceraldehyde-3p-dehydrogenase. *J. Exp. Med.* **170**, 2065–2080
41. Liska, A. J., and Shevchenko, A. (2003) Expanding the organismal scope of proteomics: cross-species protein identification by mass spectrometry and its implications. *Proteomics* **3**, 19–28
42. Juvonen, T., Parkkila, S., Parkkila, A. K., Niemela, O., Lajunen, L. H. J., Kairaluoma, M. I., Peramaki, P., and Rajaniemi, H. (1994) High-activity carbonic anhydrase isoenzyme (Ca-II) in human gallbladder epithelium. *J. Histochem. Cytochem.* **42**, 1393–1397
43. Aisen, P. (1974) Role of transferrin in iron transport. *Br. J. Haematol.* **26**, 159–163
44. Schmid, R., Diamond, I., Hammaker, L., and Gundersen, C. B. (1965) Interaction of bilirubin with albumin. *Nature* **206**, 1041
45. Lake-Bakaar, G., Dooley, J., McKavanagh, S., and Summerfield, J. A. (1979) Clearance of immunoreactive trypsin in bile. *Gut* **20**, A934–A935
46. Hanna, R. E. B. (1980) *Fasciola hepatica*: an immunofluorescent study of antigenic changes in the tegument during development in the rat and sheep. *Exp. Parasitol.* **50**, 155–170
47. Braschi, S., Curwen, R. S., Ashton, P. D., Verjovski-Almeida, S., and Wilson, A. (2006) The tegument surface membranes of the human blood parasite *Schistosoma mansoni*: a proteomic analysis after differential extraction. *Proteomics* **6**, 1471–1482
48. Donnelly, S., O'Neill, S. M., Sekiya, M., Mulcahy, G., and Dalton, J. P. (2005) Thioredoxin peroxidase secreted by *Fasciola hepatica* induces the alternative activation of macrophages. *Infect. Immun.* **73**, 166–173
49. Salazar-Calderon, M., Martin-Alonso, J. M., de Eguino, A. D. R., Casais, R., Marin, M. S., and Parra, F. (2000) *Fasciola hepatica*: heterologous expression and functional characterization of a thioredoxin peroxidase. *Exp. Parasitol.* **95**, 63–70
50. Mulcahy, G., and Dalton, J. P. (1998) Vaccines in the control of liver fluke infection in ruminants: current status and prospects. *Irish Vet. J.* **51**, 520–525
51. Braun, U., Wolfensberger, R., and Hertzberg, H. (1995) Diagnosis of liver flukes in cows—a comparison of the findings in the liver, in the feces, and in the bile. *Schweiz. Arch. Tierheilkd.* **137**, 438–444