

Definition of *Arabidopsis* Sterol-rich Membrane Microdomains by Differential Treatment with Methyl- β -cyclodextrin and Quantitative Proteomics*[§]

Sylwia Kierszniowska, Bettina Seiwert, and Waltraud X. Schulze‡

Plasma membranes are dynamic compartments with key functions in solute transport, cell shape, and communication between cells and the environment. In mammalian cells and yeast, the plasma membrane has been shown to be compartmented into so-called lipid rafts, which are defined by their resistance to treatment with non-ionic detergents. In plants, the existence of lipid rafts has been postulated, but the precise composition of this membrane compartment is still under debate. Here we were able to experimentally clearly distinguish (i) true sterol-dependent “raft proteins” and (ii) sterol-independent “non-raft” proteins and co-purifying “contaminants” in plant detergent-resistant membranes. We used quantitative proteomics techniques involving ¹⁵N metabolic labeling and specific disruption of sterol-rich membrane domains by methyl- β -cyclodextrin. Among the sterol-dependent proteins we found an over-representation of glycosylphosphatidylinositol-anchored proteins. A large fraction of these proteins has functions in cell wall anchoring. We were able to distinguish constant and variable components of plant sterol-rich membrane microdomains based on their responsiveness to the drug methyl- β -cyclodextrin. Predominantly proteins with signaling functions, such as receptor kinases, G-proteins, and calcium signaling proteins, were identified as variable members in plant lipid rafts, whereas cell wall-related proteins and specific proteins with unknown functions make up a core set of sterol-dependent plant plasma membrane proteins. This allows the plant to maintain a balance between static anchoring of cell shape forming elements and variable adjustment to changing external conditions. *Molecular & Cellular Proteomics* 8:612–623, 2009.

Since the first description of microdomains with distinct lipid composition in plasma membranes of mammalian cells (1, 2), the concept of a key role of such “lipid rafts” in cellular

processes has been proposed for a range of biological functions. These membrane microdomains are biochemically defined by their resistance to treatment with the non-ionic detergents in low temperature and can be isolated as a distinct fraction in gradient centrifugation.

A role of specific lipid domains composed of sterols and sphingolipids for cellular sorting and trafficking processes was proposed more than 20 years ago (3). It is hypothesized that the distinct lipid composition of lipid rafts creates a special environment that facilitates protein-protein interactions, protein recruitment in cellular trafficking events, and endocytosis and signaling (4–6). Natural sterols and sphingolipids from plants, fungi, or animals have been shown to be capable of inducing the formation of membrane heterogeneities (7). Recent studies on giant blebs produced from cultured mammalian cells allowed visualization of patchy lipid heterogeneity (8). These patches had a temperature-dependent size up to micrometer scales.

In mammalian cells, lipid rafts have been shown to play a role in many different events: endocytosis via caveolae (9), virus budding or pathogen entry (10), regulation of exocytosis (11), actin cytoskeleton organization (12), and apoptosis (13). Cholesterol-dependent segregation of lipid raft proteins from non-raft proteins was visualized in mammalian cells and is consistent with the view that raft domains in the plasma membrane of cells are usually small and highly dispersed, but their size can be modulated by oligomerization of raft components (2).

Although in the mammalian system and in yeast the concept of lipid rafts and their role in cellular processes has been widely studied using a variety of biochemical and cell biological tools, in plants our actual knowledge of the composition and role of such membrane microdomains remained under debate for a long time (14). In recent years, several proteomics studies were carried out on detergent-resistant membrane fractions of *Arabidopsis*, tobacco, mustard seedlings, or *Medicago* nodules (15–18). However, most of these studies present long lists of proteins identified in the biochemical preparation of detergent-resistant domains and compare them with whole plasma membrane preparations. Thus, a quantitative proteomics study allowing a distinction of proteins co-purifying in the detergent-resistant membrane

From the Max-Planck-Institut für Molekulare Pflanzenphysiologie, Am Mühlenberg 1, 14476 Golm, Germany

✂ Author's Choice—Final version full access.

Received, July 29, 2008, and in revised form, November 21, 2008

Published, MCP Papers in Press, November 25, 2008, DOI 10.1074/mcp.M800346-MCP200

(DRM)¹ fraction *versus* true proteins dependent on sterol-rich membrane regions has not yet been carried out in plants.

The aim of this study was to characterize the composition of plasma membrane detergent-resistant domains using comparative quantitative proteomics techniques in combination with selective disruption of the sterol-rich membrane domains by chemical treatment. Thereby a differentiation between co-purifying “non-raft” proteins and sterol-dependent “raft-associated” proteins was possible, giving new insights into the lipid raft model in plants.

MATERIALS AND METHODS

Arabidopsis Cell Suspension Cultures—*Arabidopsis thaliana* Col-0 cell suspension cultures (19) were grown in full medium as described previously (20). For quantitative proteomics experiments, half of the cell cultures were metabolically labeled with ¹⁵N by growth under [¹⁵N]KNO₃ as the only nitrogen source for at least two passages over 2 weeks (21).

Preparation of Plasma Membrane and Detergent-resistant Fraction—Plasma membranes were purified from the microsomal pellet (100,000 × *g*) over a two-phase system of dextran and polyethylene glycol (22) using 6.4% dextran, 6.4% polyethylene glycol, 5 mM phosphate buffer, pH 7.8, and 5 mM KCl. Protein concentration was determined by Bradford assay. The plasma membrane pellet was then treated with the non-ionic detergent Triton X-100 for 30 min at 4 °C at a ratio of protein to detergent of 1:15 and shaken at 60 rpm. Treated membranes were then taken up in 2.4 M sucrose to a final concentration of 1.8 M and overlaid by a sucrose step gradient from 1.8 to 0.15 M. After gradient centrifugation at 250,000 × *g* for 18 h, a ringlike structure was visible at the interface of 1.4 and 0.15 M sucrose concentrations. The fractions of 1 ml above and 1 ml below the interface were collected as the detergent-resistant membrane fraction. The DRM fraction was diluted five times with 10 mM Tris-HCl buffer, pH 8, and membranes were pelleted at 200,000 × *g* for 1 h.

Isolated plasma membranes (300 μg) were treated with the sterol-disrupting agent methyl-β-cyclodextrin (mβcd) at a final concentration of 5, 15, or 30 mM for 1 h at 37 °C as described previously (23). Membranes were then washed in phosphate buffer and pelleted prior to Triton X-100 treatment for DRM preparation.

Protein Analysis and Identification by LC/MS/MS—DRMs were pelleted at 100,000 × *g*. Pellets were resuspended in 6 M urea and 2 M thiourea for denaturation and were subsequently subjected to reduction in DTT, and free SH groups were carbamidomethylated using iodoacetamide (24). This was followed by digestion with Lys-C and with trypsin (24). Digested samples were desalted over C₁₈ Stop And Go Extraction tips (Empore Disk, Varian Inc.) (25).

Tryptic peptide mixtures were analyzed by LC/MS/MS using nano-flow HPLC (Proxeon Biosystems) and an Orbitrap hybrid mass spectrometer (LTQ-Orbitrap, Thermo Electron) as mass analyzer. Peptides were eluted from a 75-μm analytical column (Reprosil C₁₈, Dr. Maisch GmbH) on a linear gradient running from 4 to 64% acetonitrile in 90 min and sprayed directly into the LTQ-Orbitrap mass spectrometer. Proteins were identified by MS/MS by information-dependent acquisition of fragmentation spectra of multiple charged peptides. Up to five data-dependent MS/MS spectra were acquired in the linear ion trap for each FTMS full-scan spectrum acquired at 30,000 full-width

half-maximum resolution settings with an overall cycle time of approximately 1 s. Fragment MS/MS spectra from raw files were extracted as DTA files and then merged to peak lists using the default settings of DTASuperCharge version 1.18 (SourceForge, Inc.) with a tolerance for precursor ion detection of 50 ppm.

Fragmentation spectra were searched against a non-redundant *Arabidopsis* protein database (TAIR8, version April 2008; 31,921 entries) using the Mascot algorithm (version 2.2.0; Matrix Science). The database contained the full *Arabidopsis* proteome and commonly observed contaminants (human keratin, trypsin, and lysyl endopeptidase); thus no taxonomic restrictions were used during the automated database search. The following search parameters were applied: trypsin as cleaving enzyme; peptide mass tolerance, 10 ppm; MS/MS tolerance, 0.8 Da; and one missed cleavage allowed. Carbamidomethylation of cysteine was set as a fixed modification, and methionine oxidation was chosen as a variable modification. “¹⁵N metabolic labeling” was chosen as a quantitative method for Mascot database searching, allowing identification of labeled and unlabeled peptides within the same database search. Only peptides with a length of more than five amino acids were considered.

In general, peptides were accepted without manual interpretation if they displayed a Mascot score greater than 31 (as defined by Mascot $p < 0.01$ significance threshold). Peptides with a score greater than 24 were manually inspected requiring a series of three consecutive y or b ions to be accepted; in addition, mass accuracy and delta scores were taken into account when single peptides were accepted. Original MS/MS spectra considered for quantitation in this analysis will be deposited in the Promex spectral library, and spectra of accepted “single peptide” hits are included in supplemental Fig. 1. Using the above criteria for protein identification, the rate of false identifications as determined by the “decoy database” function implemented in Mascot version 2.2.0 was 3.4% on a 95% confidence level. The false positive rate in ¹⁵N-labeled samples was higher because of increased ambiguity in protein identification resulting from a higher number of isobaric amino acids in labeled samples (26).

Peptide assignment to proteins was done according to the Mascot default settings, *i.e.* each redundant peptide was primarily assigned to the highest scoring protein. Isoforms of protein only appear in the tables as a separate protein entry if they were assigned at least one unique peptide. A summary of all identified peptides and their respective quantitative values is presented in supplemental Table 2.

Quantitative Protein Analysis—Ratios between labeled and unlabeled forms of each tryptic peptides were calculated in MSQuant version 1.4.3 (released May 30, 2008; SourceForge, Inc.). Quantitative information was taken from extracted ion chromatograms of the labeled and unlabeled form of each identified peptide. Thereby co-elution of both peptide forms was made a requirement, and it was manually inspected in MSQuant that the pairs of labeled and unlabeled forms actually fit with the expected isotope envelope distributions. Peptides that did not meet these criteria were omitted from the analysis as described previously (21).

Intensity ratios of the labeled ¹⁵N form to unlabeled ¹⁴N form of each identified peptide were averaged across all peptides belonging to the same protein within one experimental set. Peptides conserved in multiple members of a protein family were identified using the “show subsets” option in Mascot, and the respective peptides present in multiple proteins were excluded from quantitative analysis if the redundant peptides displayed ratios significantly different ($p < 0.05$; χ^2 test) from unique peptides of the same protein (27, 28). Peptides meeting the criteria for sequence identification but for which only ¹⁴N forms or only ¹⁵N forms were quantified were manually assigned the ratios 0.1 (¹⁴N form only) or 10 (¹⁵N form only). This affected only peptides for which the pairing labeled or unlabeled peak was at noise level. Because quantitative information was extracted from full-scan

¹ The abbreviations used are: DRM, detergent-resistant membrane; mβcd, methyl-β-cyclodextrin; GPI, glycosylphosphatidylinositol; APCI, atmospheric pressure chemical ionization; UPLC, ultraperformance LC; SUBA database, *Arabidopsis* Subcellular Database; GFP, green fluorescent protein.




FIG. 1. **Work flow of the reciprocal labeling experiments.** Two experiments were carried out in parallel. In one case, the ^{14}N cells were subjected to the $m\beta cd$ treatment, and ^{15}N cells were used as control. In the second case, the ^{15}N cells were used for $m\beta cd$ treatment, whereas the ^{14}N cells were used as untreated control. In addition, 1:1 mixtures of untreated ^{14}N and ^{15}N cells were used to define inherent differences between the cell cultures and the technical variation. The complete reciprocal experimental design was repeated four times independently. *PM*, plasma membrane.

spectra with a very low level of noise as obtained in the Orbitrap mass analyzer, no minimum threshold was set for quantitation (29). Protein abundance ratios were converted into \log_2 values. Only those proteins were considered for further analysis for which intensity ratios were obtained in both paired reciprocal experiments. Ratios of ^{15}N to ^{14}N forms and the respective standard deviation as calculated by MSQuant for each identified peptide and the number of peptides used for quantitation for each protein are presented in supplemental tables. The average relative error of quantitation for all 465 quantified proteins was 10.6%.

Reciprocal Labeling Experimental Setup and Statistical Analysis—For comparative proteomics analysis, plasma membranes of ^{15}N -labeled cell cultures were treated with $m\beta cd$, whereas plasma membranes of control cells were left untreated. In a paired reciprocal experiment using the same starting material, ^{14}N cells were treated with $m\beta cd$, whereas ^{15}N cells were left untreated. Labeled and unlabeled plasma membranes were treated with Triton X-100 in one combined sample prior to DRM preparation over a sucrose gradient.

The work flow of the experimental setup is shown in Fig. 1. The paired reciprocal experiments (Fig. 1) were independently repeated four times. Protein ratios of each replicate experimental set were z-score-standardized to allow comparison between experiments. In result tables and figures, the averages and S.D. of z-transformed protein ratios from all biological replicate experiments are presented. In general, 60% of all identified proteins were found in at least two independent reciprocal experimental setups, and 25% of all identified proteins were found in all four replicate experiments (supplemental Fig. 2).

The reciprocal labeling setup was chosen over an experimental setup using the same ^{15}N -labeled cultures as repeated internal standard (30) to specifically distinguish which proteins are responding to $m\beta cd$ (treatment effects) from those proteins that are *a priori* different between labeled and unlabeled cell cultures (*i.e.* culture effects) (31).

In summary, the data analysis work flow is based on first determining the variation between cultures based on $^{15}\text{N}/^{14}\text{N}$ ratios in independent 1:1 mixtures before $m\beta cd$ treatment is applied. The

ratios in two control experiments showed normal distribution (see Fig. 2B) and were used to define ratio-dependent standard deviations (31). In a second step, the distances to the diagonal in a graphic display of ratios in reciprocal experiments (Fig. 2A) was calculated. Then for each data point the ratio between the distance and the S.D. was calculated, and the p value was determined by a two-tailed t distribution. Subsequently a multiple testing correction was applied to the whole data set using the false discovery rate method introduced by Benjamini and Hochberg (32). Reported proteins correspond to a cutoff false discovery rate of 5%.

Analysis of Sterols by APCI/MS—Lipids were extracted from equal amounts (150 μ g) of $m\beta$ cd-treated and untreated plasma membrane preparations by a Bligh and Dyer (33) method. A final amount of 10 μ g of 3 β -hydroxy-5 α -cholestane (Sigma-Aldrich) was used as internal standard. Extracted lipids were hydrolyzed with 4 M HCl for 1 h at 80 °C and extracted with 2-hexane. The solvents were evaporated, and samples were diluted in 1:1 chloroform:methanol.

Sterols were subsequently separated by UPLC (Waters) over a C₁₈ column during a 25-min gradient using water and acetonitrile as mobile phases. Eluting peaks were directly ionized by APCI before detection in an LTQ-Orbitrap mass spectrometer (Thermo Fisher). Ions were detected in the m/z range 100–1000, and 500,000 ions were collected in the Orbitrap detector set to a resolution of 30,000 full-width half-maximum. Total ion chromatograms were extracted, and peak volumes were compared relative to the internal standard 3 β -hydroxy-5 α -cholestane.

Bioinformatics Analysis and Functional Classification—Proteins were classified to functional groups using the MapMan nomenclature developed for plant genes (34). GPI-anchored proteins were annotated based on published literature and predictions (35–37). Graphs were created using SigmaPlot version 10 (Systat Software), and statistical tests were carried out using Excel Analyze-it package version 2.07.

RESULTS

This study was designed to gain new insights about plant membrane proteins dependent on a sterol-rich membrane environment (so-called “lipid raft-proteins”). The experiments make use of the sterol-disrupting agent $m\beta$ cd in an untargeted proteomics study. The aim of this work was to disrupt lipid raft-like sterol-rich regions in the plasma membrane and thereby induce a depletion of proteins dependent on a sterol-rich membrane environment. The $m\beta$ cd-induced depletion from membranes has been described as a characteristic feature for sterol-dependent proteins (38). Thus, peptide intensity ratios between untreated and treated samples will be higher for true sterol-dependent lipid raft proteins (*i.e.* these proteins will predominantly occur in untreated samples) (23). In contrast, for co-purifying contaminants the intensity ratio from treated and untreated membranes is expected to be close to unity (*i.e.* these peptides occur in untreated and treated samples in the same amounts).

Effect of $m\beta$ cd on Plant Plasma Membrane Proteins—In total, 792 proteins were identified in DRM preparations, and for 465 proteins quantitative information from both reciprocal subsets in at least one independent reciprocal experiment was obtained. About 15% of all the proteins with reciprocal quantitative data were found to be significantly depleted from DRM preparations upon $m\beta$ cd treatment of plasma mem-

branes (Fig. 2A, *blue dots*). A full list of proteins that were found to be depleted from DRM preparations upon $m\beta$ cd treatment in at least one reciprocal experiment is presented in supplemental Table 1. A core set of 37 proteins was identified as $m\beta$ cd-responsive in DRM preparations of at least two independent reciprocal experimental sets (Table I). These $m\beta$ cd-responsive proteins are primary candidates for being associated with or located in a sterol-rich membrane micro-environment (lipid rafts). Among all proteins identified as sterol-dependent proteins, we observed a significant ($p < 0.03$, Fisher exact test) over-representation of single transmembrane domain proteins (supplemental Fig. 3). We found about 2% of the quantified proteins to be enriched in DRM preparations after $m\beta$ cd treatment in individual experiments (Fig. 2A, *red dots*).

Specificity of the $m\beta$ cd Response—To test whether proteins were removed from the plasma membrane upon $m\beta$ cd treatment or whether they were rather dissipated within the membrane, $m\beta$ cd-treated purified plasma membranes were pelleted, and proteins in the supernatant were analyzed. The majority of proteins identified in supernatants after pelleting $m\beta$ cd-treated plasma membranes were cell wall-related GPI-anchored membrane proteins, remorin, and a phosphate transporter (Table I). An increase in $m\beta$ cd concentration resulted in an increased release to the supernatant of specifically those proteins that were also classified as sterol-dependent proteins in the reciprocal labeling experiments carried out on DRM preparations. The abundance ratio of these proteins in supernatants of treated *versus* untreated plasma membranes increased in the $m\beta$ cd concentration range of 5–15 mM and saturated with an $m\beta$ cd concentration of 30 mM (Fig. 3). Overall these results support our view that $m\beta$ cd removes sterol-dependent proteins from the membrane environment in a concentration-dependent manner and that proteins are removed from the membrane by $m\beta$ cd in a form that does not sediment at $250,000 \times g$.

Effects of $m\beta$ cd on Plant Plasma Membrane Sterols—Sito-sterol and campesterol were found to be the most abundant sterols in *Arabidopsis* plasma membranes as measured by UPLC/APCI/MS (Fig. 4A). Other sterols, such as cholesterol, avenasterols, stigmasterol, and episterol, were present in up to 2 orders of magnitude lower amounts. Upon treatment with $m\beta$ cd, the identified sterols were removed from DRM preparations in a concentration-dependent manner (Fig. 4B). Treatment of plasma membranes with 30 mM $m\beta$ cd for 1 h led to a reduction of membrane sterols to about 50% of the levels present in untreated DRM fractions.

Functional Classification of $m\beta$ cd-responsive Proteins—Specifically sterol-disrupting treatment of plasma membranes with $m\beta$ cd caused a clearly significant ($p < 0.01$; Fisher exact test) depletion of specific classes of proteins from DRM preparations. Fig. 5A displays the distribution of 75 $m\beta$ cd-responsive and 263 non-responsive proteins to different MapMan functional categories (34). Most (87%) of

Effect of mβcd treatment on protein abundance in DRM





FIG. 2. A, response of proteins to mβcd treatment. Log₂ values of ¹⁵N to ¹⁴N ratios from one experiment were plotted against log₂ values of ¹⁵N to ¹⁴N ratios from the reciprocal experiment. Blue and red symbols indicate those sterol-dependent and sterol-independent proteins that show significant reciprocal response. Yellow symbols indicate those proteins that do not respond to the mβcd treatment (31). B, distribution of ratios from two control experiments (yellow), mβcd-responsive proteins (blue), and other proteins (red). The mβcd-responsive proteins fall into two classes: a core set of very strongly responsive (dark blue) and some less responsive ones (light blue).



the proteins showing significant depletion upon mβcd treatments have been localized previously to the plasma membrane by mass spectrometry or by GFP fusions as noted by the SUBA database of subcellular localizations (39). Specifically a third of these responsive plasma membrane proteins

has also been annotated as extracellular protein in the SUBA database. In contrast, among the “non-responsive” proteins we found 71% as annotated membrane proteins, and only a total of 3% has been described as extracellular proteins (Fig. 5B).

TABLE 1
Core list of proteins identified as *mβcd*-responsive in at least two biological replicate experimental sets

Each protein was functionally classified by MapMan (34). *p* values indicate the significance of the *mβcd* response. A cutoff *p* value of 0.05 was used after correction for multiple testing (32). "DRM" refers to proteins identified as *mβcd*-responsive based on differential analysis of DRM preparations after *mβcd* treatment of plasma membranes. "sup" identifies *mβcd* responsiveness based on release of these proteins to the supernatant after pelleting *mβcd*-treated plasma membranes. Letters indicate proteins previously published as being GPI-anchored (gpi) (35, 36, 50, 51) or present in DRM in *Arabidopsis* (drm) (15). Localization is taken from the SUBA database of subcellular localizations (39): PM, plasma membrane; EX, extracellular space; P, plastid; V, vacuole; C, cytosol; N, nucleus; M, mitochondrion. AGI, *Arabidopsis* gene identifier; DREPP, developmentally regulated plasma membrane polypeptide; v-ATPase, vacuolar type H⁺-ATPase; LYSM, lysin motif.

AGI	Description	MapMan functional category	<i>p</i> value	Times found	Experiment	Experimental subcellular localization from SUBA		DRM/GPI
						MS	GFP	
At3g07390	AI12; extracellular matrix structural constituent	Hormone metabolism	2.06e-06	4	sup, DRM	ER, EX, P	EX, PM	gpi
At4g26690	MRH5/SHV3; glycerophosphodiester phosphodiesterase/kinase	Lipid metabolism	2.06e-06	4	sup, DRM	ER, EX, PM, M	EX, PM	gpi
At4g31140	Glycosyl hydrolase family 17 protein	Cell wall	2.06e-06	3	sup, DRM	ER, EX, PM	EX, PM	gpi
At5g58090	Glycosyl hydrolase family 17 protein	Not assigned, no ontology	2.06e-06	4	sup, DRM	ER, EX, PM	EX, PM	gpi
At5g58480 ^a	Glycosyl hydrolase family 17 protein	Not assigned, no ontology	2.06e-06	2	DRM	C, ER, EX, PM, M	EX, PM	gpi
At3g13560	Glycosyl hydrolase family 17 protein	Cell wall	2.54e-06	3	sup, DRM	ER, EX, PM, M	EX, PM	gpi
At5g44130	Fascilin-like arabinogalactan protein (FLA13)	Cell wall	7.10e-06	6	sup, DRM	ER, EX, PM, M	EX, PM	gpi
At2g25060	Plastocyanin-like domain-containing protein	Cell wall	1.29e-05	4	sup, DRM	ER, EX, PM	EX, PM	gpi
At4g25240	SKS1 (SKU5 similar 1); copper ion-binding	Cell wall	1.56e-05	6	sup, DRM	ER, EX, PM	PM	drm, gpi
At5g55480	Glycerophosphoryldiester phosphodiesterase family protein	Lipid metabolism	1.56e-05	6	sup, DRM	EX, PM, P, M	EX, PM	drm, gpi
At4g31840	Plastocyanin-like domain-containing protein	Cell wall	1.72e-05	6	sup, DRM	ER, EX, M, V	PM	gpi
At4g12730	Fascilin-like arabinogalactan protein (FLA2)	Cell wall	1.84e-05	6	sup, DRM	ER, EX, PM, M	PM	gpi
At3g02740	Aspartyl protease family protein	Protein degradation	3.06e-05	2	sup, DRM	ER, EX, V	PM	gpi
At1g64760	Glycosyl hydrolase family 17 protein	Cell wall	4.62e-05	5	sup, DRM	ER, EX, PM	PM	gpi
At5g53870	Plastocyanin-like domain-containing protein	Cell wall	5.16e-05	4	sup, DRM	ER, EX, M, N, V	PM	gpi
At2g04780	Fascilin-like arabinogalactan protein (FLA7)	Cell wall	9.75e-05	6	sup, DRM	ER, EX, PM	EX, PM	gpi
At2g45470	Fascilin-like arabinogalactan protein (FLA8)	Cell wall	2.01e-04	5	sup, DRM	ER, EX	EX, PM	gpi
At5g55730	Fascilin-like arabinogalactan protein (FLA1)	Cell wall	2.25e-04	5	sup, DRM	ER, EX, P, M	EX, PM	gpi
At3g61260	DNA-binding family protein/remorin family protein	RNA regulation of transcription	3.35e-04	2	DRM	N	PM	gpi
At3g58100 ^a	Glycosyl hydrolase family 17 protein	Cell wall	3.99e-04	2	DRM	EX	PM	gpi
At2g17120	LYM2 (LYSM domain GPI-anchored protein 2)	Not assigned, no ontology	6.53e-04	4	DRM	C, ER, EX, P	EX, PM	gpi
At5g51480	SKS2 (SKU5 similar 2); copper ion binding	Cell wall	2.20e-03	6	sup, DRM	ER, EX, PM	PM	gpi
At2g45820	DNA-binding protein, putative	RNA regulation of transcription	3.65e-03	4	sup, DRM	C, N	PM	gpi
At5g42100	Glycosyl hydrolase family 17 protein	Cell wall	4.64e-03	2	DRM	ER, EX, M	EX, PM	gpi
At3g03520 ^a	Phosphoesterase family protein	Not assigned, no ontology	6.08e-03	2	sup	EX, M, N	V	gpi
At4g12420	SKU5 (skewed 5); copper ion binding	Not assigned, no ontology	1.29e-02	2	sup	ER, EX, PM	EX, PM	drm, gpi
At5g48450	SKS3 (SKU5 Similar 3); copper ion binding	Cell wall	1.39e-02	2	DRM	ER, EX, V	PM	gpi
At5g25090	Plastocyanin-like domain-containing protein	Cell wall	1.88e-02	2	DRM	ER, EX, PM, M	EX, PM	gpi
At4g30190	AHA2 (<i>Arabidopsis</i> H ⁺ -ATPase 2); ATPase	Transport, p- and v-ATPases	1.91e-02	2	sup	C, PM	PM	gpi
At2g01210	Leucine-rich repeat transmembrane protein kinase, putative	Signaling, receptor kinases	2.05e-02	2	DRM	C, ER, EX, PM	EX, PM, P	gpi
At2g05920	Subtilase family protein	Protein degradation	2.13e-02	2	DRM	ER, EX, M, P	EX	gpi
At4g27520	Plastocyanin-like domain-containing protein	Cell wall	2.64e-02	3	sup, DRM	ER, EX, N	EX, PM, P	gpi
At1g11820	Hydrolase, hydrolyzing O-glycosyl compounds	Not assigned, no ontology	2.99e-02	2	DRM	ER, EX, PM	EX	gpi
At2g38940	ATPT2 (phosphate transporter 2)	Transport phosphate	3.23e-02	3	sup, DRM	EX, PM, P, M	PM, V	gpi
At3g08510	ATPLC2 (phospholipase C2); phospholipase C	Lipid metabolism	3.53e-02	2	DRM	C, M, P	PM	gpi
At4g36750	Quinone reductase family protein	Lipid metabolism	4.54e-02	2	DRM	C, M, N	PM	gpi
At4g20260	DREPP plasma membrane polypeptide family protein	Not assigned, no ontology	4.71e-01	4	sup, DRM	M, N	PM, P	gpi

^a Proteins identified based on a single peptide (see supplemental Fig. 1 for annotated spectra).




FIG. 3. Concentration dependence of the $m\beta cd$ -induced depletion of proteins from *Arabidopsis* plasma membrane. The graph shows the average \log_2 value (\pm S.D.) of the ^{15}N to ^{14}N abundance ratios for proteins identified in the supernatant after pelleting of treated versus untreated plasma membranes. Only proteins identified as $m\beta cd$ -responsive in reciprocal experiments have been considered ($n = 37$).

Especially proteins with functions in cell wall structure, lipid modification, vesicle transport, and stress responses and a large number of proteins with unknown functions showed the strongest amplitude in response to $m\beta cd$ treatment. In contrast, typical co-purifying proteins, such as ribosomal proteins and vacuolar ATPases, were most abundant among proteins that did not show response to $m\beta cd$ treatment. These and other proteins with no response to $m\beta cd$ are either sterol-independent DRM proteins or contaminants.

Among the $m\beta cd$ -responsive (sterol-dependent) proteins a large number of cell wall-related proteins, fasciclin-like arabinogalactan proteins and glycosyl hydrolase family proteins were identified, and most of these proteins have been shown to be GPI-anchored (35). Proteins of primary cell wall synthesis, such as CesA proteins (CesA9 At2g21770 and CesA10 At2g25540) or Cobra (At5g60920) could not be classified because of the lack of a reciprocal quantitative value. Among the non-responsive (non-sterol-dependent) cell wall proteins were mainly proteins with functions in cell wall precursor synthesis and cell wall-degrading enzymes (supplemental Table 1).

Remorins, plant-specific plasma membrane-associated proteins of unknown structure and function, have widely been used as a marker for lipid rafts in plants (16). Two members of this protein family (At2g45820 and At3g61260) have been found as $m\beta cd$ -responsive in several independent reciprocal experiments. Thus, remorins classify as sterol-dependent proteins in our $m\beta cd$ treatment experiments.

Surprisingly proteins with functions in signaling such as receptor-like kinases, calcium signaling proteins, and G-protein signaling components showed a strong distribution to

non-sterol dependent proteins (not responsive or even enriched by $m\beta cd$), and fewer proteins in these functional groups were found to be sterol-dependent ($m\beta cd$ -responsive) proteins (Fig. 5). In total, four receptor-like kinases (At1g53730, At2g25790, At3g46290, and At5g49760) were identified as members of the sterol-dependent proteins in at least one reciprocal experimental set. In contrast, 10 other receptor-like kinases (At1g25320, At1g29740, At1g73080, At3g24540, At3g28450, At4g08850, At5g46330, At5g46330, At5g65700, and At5g67200) were identified as non-sterol-dependent proteins under the growth conditions for cell cultures in this study. Among non-sterol-dependent receptor kinases, we also found the FLS2 receptor (At5g46330) and two kinases involved in brassinosteroid signaling (At3g13380 and At3g51740). Another 35 receptor-like kinases were identified in DRM fractions but could not be classified into sterol-dependent or non-sterol-dependent proteins as they lacked a reciprocal quantitative value.

Variable and Constant Components of Sterol-rich Membrane Regions—The identified proteins with signaling functions (receptor-like kinases, calcium signaling proteins, and G-protein signaling proteins) showed a larger experiment-to-experiment deviation from the mean ratio change upon $m\beta cd$ treatment than did proteins in the functional category of cell wall, lipid-modifying, and vesicle trafficking (Fig. 6). Thus, the effect of $m\beta cd$ on signaling proteins is significantly more variable for the different proteins within a functional category and independent experimental sets compared with the effect of $m\beta cd$ on cell wall or lipid-modifying proteins. This is a strong indication for different classes of $m\beta cd$ -responsive (*i.e.* sterol-dependent) proteins. We propose that the variability of the $m\beta cd$ response among signaling proteins is an indication for a more variable and condition-dependent association of signaling proteins with sterol-rich membrane microdomains.

DISCUSSION

We used a quantitative proteomics approach to thoroughly characterize sterol-dependent membrane proteins in plant plasma membrane preparations. Our work is the first study on plant DRM that has been carried out up to date allowing such clear differentiation of sterol-dependent and sterol-independent membrane proteins. In addition it reveals novel insights into the concept of the lipid raft model in plants as a dynamic compartment.

Action of $m\beta cd$ on Plant Plasma Membranes—The precise effect of $m\beta cd$ on sterols in the plant has not been studied up to date. Therefore, it is well possible that in our experiments the plant lipid rafts were not completely disrupted by the $m\beta cd$ treatment as $m\beta cd$ may not have such a strong effect on plant-specific sterols compared with cholesterol as the dominant sterol in mammalian plasma membranes. It has been shown recently that $m\beta cd$ acts with different efficiencies on different membrane types, requiring vastly different concentrations and incubation times for different mammalian cell




FIG. 4. The effect of mβcd treatment on sterol composition of *Arabidopsis* plasma membrane. *A*, abundance of different sterols relative to the standard 3β-hydroxy-5α-cholestane. *B*, relative changes in sterol content of plasma membrane upon treatment with mβcd at 15 or 30 mM for 1 h. Averages ± S.D. of three biological replicates are shown. *d5avenasterol*, Δ⁵-avenasterol; *d7avenasterol*, Δ⁷-avenasterol.

types and tissues (40). It has also been shown on mammalian cells that mβcd removes sterol from the membranes without destroying the bilayer (41).

Our analysis of sterol composition in the DRM preparations after mβcd treatment of plasma membranes indicates that sterols are indeed being removed under the conditions applied in this study. We still observed a strong increase in the removal of sterols when increasing the mβcd concentration from 15 to 30 mM, but this effect was different for individual sterol species. However, we observed a saturation effect on the efficiency with which proteins are being removed from the plasma membrane by mβcd at concentrations between 15 and 30 mM. The mβcd concentration used for the comparative

analysis was 25 mM. Thus, it is likely that the mβcd effect on protein was saturated in this study.

Typical contaminants and co-purifying proteins in plasma membrane and DRM preparations such as ribosomal proteins, vacuolar ATPase subunits, or transcription initiation factors (15, 16) did not show any differential response to mβcd treatment. This indicates that the strategy applied here to use depletion of proteins from plasma membranes and DRM by mβcd treatment is a valid means to dissect and characterize the biochemical properties of DRM also in plants. This work is the first detailed global analysis specifically of sterol-rich membrane microdomains in plants, and by its design in reciprocal experiments it allows the precise





FIG. 5. A, distribution of mβcd-responsive proteins (blue) and proteins without an mβcd response (red) to different functional categories. Proteins were classified according to the MapMan classification scheme for plant proteins (34). n = 74 for sterol-dependent proteins and n = 295 for non-responsive proteins. B, subcellular localization from mass spectrometry and/or GFP localization of proteins identified in the mβcd disruption experiments. Subcellular localization was obtained from the SUBA database. RLKs, receptor-like kinases; ABC, ATP-binding cassette; cw, cell wall.

definition of different categories of proteins in this compartment.

The combination of the analysis of proteins removed from detergent-resistant domain preparations in reciprocal labeling

experiments with the subsequent demonstration that these particular proteins are being enriched in the supernatant after pelleting of mβcd-treated membranes is the first evidence in plants that biochemical preparations of detergent-resistant

FIG. 6. A box plot of the deviations from the mean ratio change upon $m\beta cd$ treatment for each protein in the functional categories “cell wall-related,” “lipid-modifying,” and “vesicle trafficking” as well as for signaling proteins in “receptor kinases,” “calcium signaling,” and “G-protein signaling” across four independent reciprocal experiments is shown. Proteins were classified according to the MapMan classification scheme for plant proteins (34).



domains and the actual sterol-rich membrane domains (lipid rafts) are not equal. This is an important conclusion drawn from our study, and it is well supported by work from the animal and yeast models where recent findings have strongly suggested that DRMs in standard biochemical preparations are not of the same composition as the *in vivo* functional lipid rafts in cells (42).

Characterization of Sterol-dependent Proteins—In our study, cell wall-related proteins as well as lipid- and glucan-modifying proteins were the predominant classes of proteins identified as sterol-dependent in all experiments. In contrast, proteomics studies on mammalian cell cultures using $m\beta cd$ as a cholesterol-disrupting agent revealed a strong enrichment of signaling proteins, kinases, receptors, and transporters in lipid rafts (23). Although the functional groups of proteins removed by $m\beta cd$ treatment differ between the animal and plant cells, they share common biochemistry: most proteins depleted after treatment of plant cells are experimentally proven to be GPI-anchored proteins (35, 36). Of the 37 core proteins found as responding to $m\beta cd$ in plant DRM (Table I), only eight are not modified with a glycosylphosphatidylinositol group, and one of them is predicted to be farnesylated. Furthermore 22 of the GPI-anchored proteins were also found to be released to the supernatant after pelleting of $m\beta cd$ -treated plasma membrane, and they showed the strongest amplitude of $m\beta cd$ responses in all experiments. These findings in plants are supported by earlier work on mammalian cells that the GPI anchor may be one of the (important) signals targeting a protein to lipid rafts and that the interaction of these proteins is specific to sterol-rich membrane domains (43–45). Further-

more in agreement with our findings, also quite a large percentage of proteins highly enriched as part of lipid rafts in HeLa cells (23) or human embryonic kidney cells (38) are actually GPI-anchored proteins.

The plant-specific structure of the cell wall requires anchoring and coordinated points of synthesis and modification. In fact, it has clearly been shown in plants that correct membrane targeting of GPI-anchored proteins is essential for cell wall biosynthesis. Mutants with deficient synthesis of GPI anchors display abnormal cell shapes (46). Interestingly in mammalian characterizations of lipid rafts using $m\beta cd$, many cytoskeletal proteins were found to be enriched in sterol-rich membrane domains. Thus, possibly in plants the cell wall proteins, which make up the biggest functional group of proteins identified as responding to $m\beta cd$, anchor the “plant skeleton” (cell wall) to the plasma membrane in a similar way as mammalian cells anchor their cytoskeleton (47, 48).

The Concept of Constant and Variable Proteins in Sterol-rich Membrane Regions—Surprisingly a large proportion of proteins published as “lipid raft” proteins as concluded from their proteomics detection in DRM fractions (15, 16), such as many of the receptor-like kinases, did not show drastic depletion from DRM upon $m\beta cd$ treatment. Moreover we observed an almost equal distribution of signaling proteins to sterol-dependent proteins and not sterol-dependent proteins (Fig. 3). Thus, in our experiments a substantial number of these signaling proteins did not classify as sterol-dependent proteins in our study.

We therefore conclude that sterol-rich membrane regions in plants consist of a “constant” core component mainly con-

taining cell wall-related proteins and lipid-modifying activities and a dynamic component that includes the receptor-like kinases and other signaling proteins. For example, a given receptor-like kinase may be found responsive to $m\beta cd$ in one experimental set if under these particular conditions it is associated with sterol-rich membrane regions, whereas it may not be classified as $m\beta cd$ -responsive in the experimental set carried out a few weeks later. Dynamic polarized protein localization has been observed previously in response to pathogen infections (49). When cell cultures are treated with the elicitor peptide flg22, receptor kinase FLS2 displays a strong enrichment in DRM fractions, supporting the concept of stimulation-induced membrane microdomain localization of signaling proteins.² Thus, diurnal effects of different harvesting times or subtle differences in cell culture conditions may lead to the observed experiment-to-experiment variability in sterol-dependent behavior especially among signaling proteins. The more variable $m\beta cd$ -responsive proteins such as receptor kinases and transporters may be viewed as condition-dependent raft-associated proteins that may under particular circumstances (e.g. stimulation of any kind) display $m\beta cd$ -sensitive behavior.

In contrast, cell wall proteins, lipid-modifying proteins, and a set of proteins involved in vesicle transport seem to be a more constant component of plant sterol-rich membrane microdomains independent of external conditions. These findings are in agreement with the model of rafts consisting of a core of $m\beta cd$ -sensitive proteins, most of which are GPI-anchored, and a peripheral zone with other proteins (38). The majority of proteins we identified in detergent-resistant domains were not responsive to $m\beta cd$ and thus in our view are not part of the raft model but part of membrane region that is still resistant to Triton X-100 treatment (38).

Conclusion—In summary, our study is the first comprehensive analysis of plant sterol-rich microdomains making use of the sterol-disrupting drug $m\beta cd$, which has already been used in other organisms to characterize and develop models of lipid rafts. We conclusively showed that DRM preparations in plants are not equal to sterol-rich microdomains known as lipid rafts. Rather DRMs include sterol-rich membrane subdomains but also consist of a large fraction of sterol-independent proteins not affected by $m\beta cd$ treatment. In addition, this work provides the first evidence from plants that the sterol-rich membrane domains are dynamic structures consisting of a core set of structural cell wall-related proteins with GPI anchors and a condition-dependent variable protein component mainly with signaling functions.

Acknowledgments—We thank Dirk Hincha and Lothar Willmither for stimulating discussions and critical comments.

* This work was funded by an Emmy-Noether Fellowship of the German Research Foundation (Deutsche Forschungsgemeinschaft) (to W. S.).

² N. Zappel and R. Panstruga, personal communication.

§ The on-line version of this article (available at <http://www.mcponline.org>) contains supplemental material.

‡ To whom correspondence should be addressed. Tel.: 493315678113; Fax: 493315678403; E-mail: wschulze@mpimp-golm.mpg.de.

REFERENCES

1. Simons, K., and Ikonen, E. (1997) Functional rafts in cell membranes. *Nature* **387**, 569–572
2. Harder, T., Scheiffele, P., Verkade, P., and Simons, K. (1998) Lipid domain structure of the plasma membrane revealed by patching of membrane components. *J. Cell Biol.* **141**, 929–942
3. Karnovsky, M. J., Kleinfeld, A. M., Hoover, R. L., and Klausner, R. D. (1982) The concept of lipid domains in membranes. *J. Cell Biol.* **94**, 1–6
4. Brown, D. A., and London, E. (2000) Structure and function of sphingolipid- and cholesterol-rich membrane rafts. *J. Biol. Chem.* **275**, 17221–17224
5. Simons, K., and Toomre, D. (2000) Lipid rafts and signal transduction. *Nat. Rev. Mol. Cell Biol.* **1**, 31–39
6. Bickel, P. E. (2002) Lipid rafts and insulin signaling. *Am. J. Physiol.* **282**, E1–E10
7. Xu, X., and London, E. (2000) The effect of sterol structure on membrane lipid domains reveals how cholesterol can induce lipid domain formation. *Biochemistry* **39**, 843–849
8. Baumgart, T., Hammond, A. T., Sengupta, P., Hess, S. T., Holowka, D. A., Baird, B. A., and Webb, W. W. (2007) Large-scale fluid/fluid phase separation of proteins and lipids in giant plasma membrane vesicles. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3165–3170
9. Rothberg, K. G., Ying, Y. S., Kohlhouse, J. F., Kamen, B. A., and Anderson, R. G. (1990) The glycopospholipid-linked folate receptor internalizes folate without entering the clathrin-coated pit endocytotic pathway. *J. Cell Biol.* **110**, 637–649
10. Fittipaldi, A., Ferrari, A., Zoppe, M., Arcangeli, C., Pellegrini, V., Beltram, F., and Giacca, M. (2003) Cell membrane lipid rafts mediate caveolar endocytosis of HIV-1 Tat fusion proteins. *J. Biol. Chem.* **278**, 34141–34149
11. Salaün, C., James, D. J., and Chamberlain, L. H. (2004) Lipid rafts and the regulation of exocytosis. *Traffic* **5**, 255–264
12. Falk, J., Thoumine, O., Dequidt, C., Choquet, D., and Faivre-Sarrailh, C. (2004) NrCAM coupling to the cytoskeleton depends on multiple protein domains and partitioning into lipid rafts. *Mol. Biol. Cell.* **15**, 4695–4709
13. Ayllon, V., Fleischer, A., Cayla, X., Garcia, A., and Rebollo, A. (2002) Segregation of Bad from lipid rafts is implicated in the induction of apoptosis. *J. Immunol.* **168**, 3387–3393
14. Bhat, R. A., and Panstruga, R. (2005) Lipid rafts in plants. *Planta* **223**, 5–19
15. Borner, G. H. H., Sherrier, D. J., Weimar, T., Michaelson, L. V., Hawkins, N. D., MacAskill, A., Napier, J. A., Beale, M. H., Lilley, K. S., and Dupree, P. (2005) Analysis of detergent-resistant membranes in *Arabidopsis*. Evidence for plasma membrane lipid rafts. *Plant Physiol.* **137**, 104–116
16. Mongrand, S., Morel, J., Laroche, J., Caverol, S., Carde, J.-P., Hartmann, M.-A., Bonneau, M., Somon-Plas, F., Lessire, R., and Bessoule, J.-J. (2004) Lipid rafts in higher plant cells. Purification and characterization of Triton X-100-insoluble microdomains from tobacco plasma membrane. *J. Biol. Chem.* **279**, 36277–36286
17. Morel, J., Claverol, S., Mongrand, S., Furt, F., Fromentin, J., Bessoule, J. J., Blein, J. P., and Simon-Plas, F. (2006) Proteomics of plant detergent resistant membranes. *Mol. Cell. Proteomics* **5**, 1396–1411
18. Lefebvre, B., Furt, F., Hartmann, M. A., Michaelson, L. V., Carde, J. P., Sargueil-Boiron, F., Rossignol, M., Napier, J. A., Cullimore, J., Bessoule, J. J., and Mongrand, S. (2007) Characterization of lipid rafts from Medicago truncatula root plasma membranes: a proteomic study reveals the presence of a raft-associated redox system. *Plant Physiol.* **144**, 403–418
19. Pauly, M., Eberhard, S., Albersheim, P., Darvill, A., and York, W. S. (2001) Effects of the *mur1* mutation on xyloglucans produced by suspension-cultured *Arabidopsis thaliana* cells. *Planta* **214**, 67–74
20. Jouanneau, J. P., and Peaud-Lenoel, C. (1967) Growth and synthesis of proteins in cell suspensions of a kinetin dependent tobacco. *Physiol. Plant* **20**, 834–850
21. Engelsberger, W. R., Erban, A., Kopka, J., and Schulze, W. X. (2006) Metabolic labeling of plant cell cultures with $K^{15}NO_3$ as a tool for quantitative analysis of proteins and metabolites. *Plant Methods* **2**, 1–11
22. Marmagne, A., Salvi, D., Rolland, N., Ephritikhine, G., Joyard, J., and Barbier-Brygoo, H. (2006) Purification and fractionation of membranes

- for proteomic analyses. *Methods Mol. Biol.* **323**, 403–420
23. Foster, L. J., de Hoog, C., and Mann, M. (2003) Unbiased quantitative proteomics of lipid rafts reveals high specificity for signaling factors. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 5813–5818
 24. Olsen, J. V., Ong, S.-E., and Mann, M. (2004) Trypsin cleaves exclusively C-terminal to arginine and lysine residues. *Mol. Cell. Proteomics* **3**, 608–614
 25. Ishihama, Y., Rappsilber, J., and Mann, M. (2006) Modular Stop And Go Extraction Tips with stacked disks for parallel and multidimensional peptide fractionation in proteomics. *J. Proteome Res.* **5**, 988–994
 26. Nelson, C. J., Huttlin, E. L., Hegeman, A. D., Harms, A. C., and Sussman, M. R. (2007) Implications of ¹⁵N-metabolic labeling for automated peptide identification in Arabidopsis thaliana. *Proteomics* **8**, 1279–1292
 27. Andersen, J. R., Lyon, C. E., Fox, A. H., Leung, A. K. L., Lam, Y. W., Steen, H., Mann, M., and Lamond, A. I. (2002) Direct proteomic analysis of the human nucleolus. *Curr. Biol.* **12**, 1–11
 28. Andersen, J. S., Lam, Y. W., Leung, A. K. L., Ong, S.-E., Lyon, C. E., Lamond, A. I., and Mann, M. (2005) Nucleolar proteome dynamics. *Nature* **433**, 77–83
 29. Venable, J. D., Wohlschlegel, J., McClatchy, D. B., Park, S. K., and Yates, J. R., III (2007) Relative quantification of stable isotope labeled peptides using a linear ion trap-Orbitrap hybrid mass spectrometer. *Anal. Chem.* **79**, 3056–3064
 30. Birkemeyer, C., Lüdemann, A., Wagner, C., Erban, A., and Kopka, J. (2005) Metabolome analysis: the potential of in vivo labeling with stable isotopes for metabolite profiling. *Trends Biotechnol.* **23**, 28–33
 31. Kierszniewska, S., Walther, D., and Schulze, W. (2009) Ratio-dependent significance thresholds in reciprocal ¹⁵N-labeling experiments as a robust tool in detection candidate proteins responding to biological treatment. *Proteomics*, in press
 32. Benjamini, Y., and Hochberg, Y. (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc.* **57**, 289–300
 33. Bligh, E. G., and Dyer, W. J. (1959) A rapid method for total lipid extraction and purification. *Can. J. Biochem. Physiol.* **31**, 911–917
 34. Thimm, O., Blasing, O., Gibon, Y., Nagel, A., Meyer, S., Kruger, P., Selbig, J., Muller, L. A., Rhee, S. Y., and Stitt, M. (2004) MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant J.* **37**, 914–939
 35. Elortza, F., Nühse, T. S., Foster, L. J., Stensballe, A., Peck, S. C., and Jensen, O. N. (2003) Proteomic analysis of glycosylphosphatidylinositol-anchored membrane proteins. *Mol. Cell. Proteomics* **2**, 1261–1270
 36. Elortza, F., Mohammed, S., Bunkenborg, J., Foster, L. J., Nühse, T. S., Brodbeck, U., Peck, S. C., and Jensen, O. N. (2006) Modification-specific proteomics of plasma membrane proteins: identification and characterization of glycosylphosphatidylinositol-anchored proteins released upon phospholipase D treatment. *J. Proteome Res.* **5**, 935–943
 37. Marmagne, A., Ferro, M., Meinnel, T., Bruley, C., Kuhn, L., Garin, J., Barbier-Brygoo, H., and Ephritikhine, G. (2007) A high content in lipid-modified peripheral proteins and integral receptor kinases features in the Arabidopsis plasma membrane proteome. *Mol. Cell. Proteomics* **6**, 1980–1996
 38. Ilangumaran, S., and Hoessli, D. C. (1998) Effects of cholesterol depletion by cyclodextrin on the sphingolipid microdomains of the plasma membrane. *Biochem. J.* **335**, 433–440
 39. Heazlewood, J. L., Verboom, R. E., Tonti-Filippini, J., Small, I., and Millar, A. H. (2007) SUBA: the Arabidopsis Subcellular Database. *Nucleic Acids Res.* **35**, D213–D218
 40. Zidovetzki, R., and Levitan, I. (2007) Use of cyclodextrins to manipulate plasma membrane cholesterol content: evidence, misconceptions and control strategies. *Biochim. Biophys. Acta* **1768**, 1311–1324
 41. Ohtani, Y., Irie, T., Uekama, K., Fukunaga, K., and Pitha, J. (1989) Differential effects of α -, β - and γ -cyclodextrins on human erythrocytes. *Eur. J. Biochem.* **186**, 17–22
 42. Brown, D. A. (2006) Lipid rafts, detergent-resistant membranes, and raft targeting signals. *Physiology* **21**, 430–439
 43. Cordy, J. M., Hussain, I., Dingwall, C., Hooper, N. M., and Turner, A. J. (2003) Exclusively targeting β -secretase to lipid rafts by GPI-anchor addition up-regulates β -site processing of the amyloid precursor protein. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 11735–11740
 44. Brown, D. A., and Rose, J. K. (1992) Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical cell surface. *Cell* **68**, 533–544
 45. Legler, D. F., Doucey, M. A., Schneider, P., Chapatte, L., Bender, F. C., and Bron, C. (2005) Differential insertion of GPI-anchored GFPs into lipid rafts of live cells. *FASEB J.* **19**, 73–75
 46. Gillmor, C. S., Lukowitz, W., Brininstool, G., Sedbrook, J. C., Hamann, T., Poindexter, P., and Somerville, C. (2005) Glycosylphosphatidylinositol-anchored proteins are required for cell wall synthesis and morphogenesis in Arabidopsis. *Plant Cell* **17**, 1128–1140
 47. Kwik, J., Boyle, S., Fooksman, D., Margolis, L., Sheetz, M. P., and Edidin, M. (2003) Membrane cholesterol, lateral mobility, and the phosphatidylinositol 4,5-bisphosphate-dependent organization of cell actin. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 13964–13969
 48. Palazzo, A. F., Eng, C. H., Schlaepfer, D. D., Marcantonio, E. E., and Gundersen, G. G. (2004) Localized stabilization of microtubules by integrin- and FAK-facilitated Rho signaling. *Science* **303**, 836–839
 49. Bhat, R. A., Miklis, M., Schmelzer, E., Schulze-Lefert, P., and Panstruga, R. (2005) Recruitment and interaction dynamics of plant penetration resistance components in a plasma membrane microdomain. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 3135–3140
 50. Deng, Z., Zhang, X., Tang, W., Oses-Prieto, J. A., Suzuki, N., Gendron, J. M., Chen, H., Guan, S., Chalkley, R. J., Peterman, K. T., Burlingame, A. L., and Wang, Z. Y. (2007) A proteomic study of brassinosteroid response in Arabidopsis. *Mol. Cell. Proteomics* **6**, 2058–2071
 51. Borner, G. H., Lilley, K. S., Stevens, T. J., and Dupree, P. (2003) Identification of glycosylphosphatidylinositol-anchored proteins in Arabidopsis. A proteomic and genomic analysis. *Plant Physiol.* **132**, 568–577