

Endogenous Processing and Presentation of T-cell Epitopes from *Chlamydia trachomatis* with Relevance in HLA-B27-associated Reactive Arthritis*[§]

Juan J. Cragolini, Noel García-Medel, and José A. López de Castro[‡]

Chlamydia trachomatis triggers reactive arthritis, a spondyloarthropathy linked to the human major histocompatibility complex molecule HLA-B27, through an unknown mechanism that might involve molecular mimicry between chlamydial and self-derived HLA-B27 ligands. *Chlamydia*-specific CD8⁺ T-cells are found in reactive arthritis patients, but the immunogenic epitopes are unknown. A previous screening of the chlamydial genome for putative HLA-B27 ligands predicted multiple peptides that were recognized *in vitro* by CD8⁺ T-lymphocytes from patients. Here stable transfectants expressing bacterial fusion proteins in human cells were generated to investigate the endogenous processing and presentation by HLA-B27 of two such epitopes through comparative immunoproteomics of HLA-B27-bound peptide repertoires. A predicted T-cell epitope, from the CT610 gene product, was presented by HLA-B27. This is, to our knowledge, the first endogenously processed epitope involved in HLA-B27-restricted responses against *C. trachomatis* in reactive arthritis. A second predicted epitope, from the CT634 gene product, was not detected. Instead a non-predicted nonamer from the same protein was identified. Both bacterial peptides showed very high homology with human sequences containing the HLA-B27 binding motif. Thus, expression and intracellular processing of chlamydial proteins into human cells allowed us to identify two bacterial HLA-B27 ligands, including the first endogenous T-cell epitope from *C. trachomatis* involved in spondyloarthropathy. That human proteins contain sequences mimicking chlamydial T-cell epitopes suggests a basis for an autoimmune component of *Chlamydia*-induced HLA-B27-associated disease. *Molecular & Cellular Proteomics* 8:1850–1859, 2009.

Chlamydia trachomatis is an obligate intracellular parasite that infects the urogenital epithelium. It is a very common

pathogen and one frequently inducing reactive arthritis (ReA)¹ (1). Multiple strategies, including down-regulation of major histocompatibility complex (MHC) class I and class II expression (2–4) and persistence, have been developed by the bacteria to evade the immune system. Yet both CD4⁺ and CD8⁺ T-cell responses are activated upon infection (5). In particular, HLA-B27-restricted CD8⁺ T-lymphocytes are found in patients with *Chlamydia*-induced ReA (6, 7). The role of these cells in the pathogenesis and evolution of ReA to chronic disease is probably mediated by IFN- γ . Secretion of this cytokine is critical for the protective immunity against *Chlamydia* (8) because it inhibits the bacterial growth (9). However, this is often insufficient to promote complete clearance of *C. trachomatis*, and actually IFN- γ -induced depletion of the tryptophan pool induces the differentiation of the metabolically active reticular bodies to persistent forms (10), which sustain chronic infection and ReA. The high prevalence of HLA-B27 among patients with *Chlamydia*-induced ReA (11), especially in its chronic form, suggests a pathogenetic mechanism based on interactive effects of the bacteria and HLA-B27 that seems unrelated to the capacity of *C. trachomatis* to infect or replicate into HLA-B27-positive cells (12). One such mechanism could be T-cell-mediated autoimmunity elicited by molecular/antigenic mimicry between chlamydial and self-derived HLA-B27 ligands. Antigenic mimicry between chlamydial and homologous α -myosin-derived peptides is crucial to inducing autoimmune myocarditis in mice (13). Breakdown of cytotoxic T-lymphocyte (CTL) tolerance to HLA-B27 was observed in transgenic rats upon exposure to *C. trachomatis* (14). Cross-reactivity between HLA-B27-restricted *Chlamydia*-specific CTL with self-derived HLA-B27 epitopes has not been reported. However, a biochemical basis for it was provided by the finding of an endogenously processed and presented peptide from the DNA primase of

From the Centro de Biología Molecular Severo Ochoa (Consejo Superior de Investigaciones Científicas and Universidad Autónoma de Madrid), Universidad Autónoma, 28049 Madrid, Spain

Received, February 25, 2009, and in revised form, May 6, 2009
Published, MCP Papers in Press, May 13, 2009, DOI 10.1074/mcp.M900107-MCP200

¹ The abbreviations used are: ReA, reactive arthritis; MHC, major histocompatibility complex; CTL, cytotoxic T-lymphocyte; GFP, green fluorescent protein; NQRA, Na⁺-translocating NADH-quinone reductase subunit A; PqqC, pyrroloquinoline-quinone synthase-like protein; htMDM, high throughput mass data manager; s/n, signal-to-noise ratio; IFN, interferon; fr., fraction; E3, ubiquitin-protein isopeptide ligase; TCR, T-cell antigen receptor.

C. trachomatis with high homology to a self-derived HLA-B27 ligand (15, 16).

Because of the likely involvement of HLA-B27 in the pathogenesis of chronically evolving ReA, the role of CD8⁺ T-cell responses in the protective immunity against *C. trachomatis* and the presence of HLA-B27-restricted T-cells in patients with *Chlamydia*-induced ReA, the identification of relevant chlamydial epitopes becomes crucial to establish the pathogenetic mechanism of this disease. Unfortunately a direct analysis of chlamydial HLA-B27 ligands expressed on infected cells is exceedingly difficult because of their extremely low amounts, which challenge even the most sensitive techniques of MS. In the case of *Chlamydia*, the situation is further complicated by the down-regulation of MHC class I expression shortly after infection (3, 4). To our knowledge, only one MHC class I ligand was recently identified, in the mouse system, from *Chlamydia muridarum*-infected cells using state-of-the-art MS techniques (17). Due in part to this difficulty, alternative approaches, such as expression cloning and synthetic peptide epitope mapping (18, 19) or MHC class I tetramer arrays (20), have been used to identify MHC class I-restricted chlamydial T-cell epitopes in mice. In a previous study (6) predictive algorithms were used to screen the whole genome of *C. trachomatis* for nonamer peptide sequences containing the HLA-B*2705 binding motif and a high probability of being generated by proteasomal cleavage. This led to identifying multiple sequences that, when used as synthetic peptides *in vitro*, stimulated CD8⁺ T-cells from patients with *Chlamydia*-induced ReA. Such cells could also be detected in the synovial fluid of these patients using HLA-B27 tetramers complexed to some of these peptides (7).

Although these strategies identify chlamydial sequences that are recognized by CD8⁺ T-cells they do not prove that these peptides are the endogenously processed epitopes that activated the natural T-cell responses to the bacteria *in vivo*. Because of the intrinsic cross-reactivity of T-cells (21, 22), it is conceivable that synthetic peptides recognized *in vitro* may be different from the natural epitopes generated by endogenous processing of the chlamydial proteins that elicit the HLA-B27-restricted T-cell responses in ReA patients. To investigate this issue we focused on two predicted epitopes (6). Stable transfectants expressing the corresponding chlamydial proteins fused to green fluorescent protein (GFP) were generated in a B*2705-positive cell line. The endogenous processing and presentation of the predicted epitopes or other peptides from the same bacterial protein were analyzed by comparative immunoproteomics analysis of the B*2705-bound peptide repertoires from transfected and untransfected cells and sequencing of peptides differentially presented on the bacterial protein transfectant.

EXPERIMENTAL PROCEDURES

Bacterial Gene Constructs and Transfectants—GFP bacterial fusion proteins were produced fusing the cDNA coding for the Na⁺-

translocating NADH-quinone reductase subunit A (NQR) or pyrrolo-quinoline-quinone synthase-like protein (PqqC) (CT634 and CT610 gene products, respectively) of *C. trachomatis* serovar L2 (Advanced Biotechnologies, Columbia, MD) to the 3'-end of the GFP gene. Both cDNAs were amplified by PCR using the following primers: 5'-TCTCTCTCGAATTCTATGAAATAGTTGTTTCTCGCGGA and 3'-TCTCTCTCGGATCCTAACGAGGAGGTTACCACATT for NQR and 5'-CTCTCTCTAGATCTATGATGGAGGTGTTTATGAATTTT and 3'-CTCTCTCTCGACATAAGATTGATGACAACACTACAACA for PqqC. The complete sequences were cloned into the pGFP-C1 vector (BD Biosciences Clontech) in the EcoRI and BamHI sites and in the BglII and Sall sites, respectively.


Stable transfectants were generated and characterized as previously described (16). Briefly the human HLA-A,B-deficient lymphoid cell line C1R (23) was electroporated with the GFP or GFP-chlamydial gene constructs and the RSV5 vector carrying the hygromycin resistance gene. Fluorescent hygromycin-resistant cells were selected by flow cytometry, and the expression of the fusion protein was analyzed by Western blot.

Isolation of HLA-B27-bound Peptides—B*2705-bound peptides were isolated from $1-2 \times 10^{10}$ C1R-B*2705 cells and C1R-B*2705 transfectants expressing chlamydial proteins as described previously (24). Briefly cells were lysed in 1% Igepal CA-630 (Sigma), 20 mM Tris/HCl buffer, 150 mM NaCl, pH 7.5 in the presence of a mixture of protease inhibitors. HLA-B27-peptide complexes were isolated by affinity chromatography of the soluble fraction with the W6/32 monoclonal antibody (IgG2a, specific for a monomorphic HLA-A,B,C determinant) (25). HLA-B27-bound peptides were eluted at room temperature with 0.1% aqueous TFA, concentrated with Centricon 3 (Amicon, Beverly, MA), and fractionated by HPLC as described previously (26).

MALDI-TOF MS Analysis—HPLC fractions were analyzed in a MALDI-TOF/TOF mass spectrometer (4800 Proteomics Analyzer, Applied Biosystems, Foster City, CA). The samples were dried down with a SpeedVac system (Savant Global Medical Instrumentation, Ramsey, MN), reconstituted in 0.6 μ l of TA buffer (33% aqueous acetonitrile, 0.1% TFA), loaded onto an Opti-TOF™ 384-well MALDI insert (Applied Biosystems), and allowed to dry at room temperature. Then 0.6 μ l of matrix solution (α -cyano-4-hydroxycinnamic acid (Bruker Daltonics, Bremen, Germany)) at 3 mg/ml was added and allowed to dry at room temperature. MS data were acquired in the mass range 800–2000 Da in reflector positive mode at 25 kV and analyzed using the Data Explorer software, version 4.9 (Applied Biosystems). Each spectrum was externally calibrated with the Peptide Calibration Standard Mixture (Bruker Daltonics, product number 206195) to reach a typical mass measurement accuracy of <25 ppm.

Automated Comparative Analysis of MALDI-TOF MS Data—A systematic comparison of the MALDI-TOF MS spectra of the HPLC fractions from HLA-B27-bound peptide pools was carried out with an algorithm programmed in Visual Basic 6.0 and implemented as a macro within Microsoft® Excel, designated as high throughput mass data manager (htMDM). This tool recognizes the raw numeric data list from each individual mass spectrum from the MALDI-TOF/TOF mass spectrometer software. First, for comparison of distinct peptide pools, the ion peaks with a signal-to-noise ratio (s/n) smaller than 3-fold, relative to the average s/n of the MS spectrum, were removed. Second, ion peaks with a s/n below 5% of the maximal s/n observed in the MS spectrum were also ignored. Third, an exclusion list of matrix-related ion peaks was generated from direct MS analysis of the matrix alone. The ion peaks whose *m/z* was identical (± 0.4) to those in this list were excluded. For comparison of MS spectra corresponding to correlative (± 1) HPLC fractions from two peptide pools, ion peaks were considered to reflect identical peptides when their observed *m/z* were within a user-defined range, which depends on the

FIG. 1. Summary of the strategy used to identify endogenously processed HLA-B27 ligands from *C. trachomatis*. The B*2705-bound peptide repertoires were isolated in parallel experiments from about $1\text{--}2 \times 10^{10}$ C1R transfectant cells expressing (2) or not expressing (1) a GFP bacterial fusion protein by immunopurification of HLA-B27 and acid extraction (3). The peptide pools were fractionated by HPLC under identical conditions and in consecutive runs (4). Each individual HPLC fraction was analyzed by MALDI-TOF MS, and the corresponding MS spectra of correlative fractions were systematically compared to identify peptides found only in the bacterial protein transfectant (5). The amino acid sequence of the differentially expressed peptides was determined by MS/MS fragmentation analysis (6). The corresponding peptide was chemically synthesized, and the MS/MS spectrum of the synthetic peptide was compared with that of the endogenous ligand (7).



accuracy of the instrument used. In our study this range was set at ± 0.4 .

The htMDM tool also allows searching for specific ion peaks, such as candidate peptide epitopes or peptides of a particular size. A user-defined inclusion list with the m/z values of interest can be stored and used to automatically screen all the MS spectra for the presence of ion peaks with identical (± 0.4) m/z for their quick identification. The htMDM output consists of a listing of filtered m/z values with the corresponding retention time or HPLC fraction number. Matches with the exclusion and inclusion directories are marked as well as shared and specific m/z signals from each of the peptide repertoires compared. To verify the performance of the htMDM tool, both automatic and manual comparisons were carried out.

Peptide Sequencing—MALDI-TOF-TOF MS/MS fragmentation spectra were acquired with Data Explorer, version 4.9, at 1 kV using collision-induced dissociation with atmospheric air and a precursor mass window of ± 2.5 Da. A s/n of 10 was used for processing data. Interpretation of the MS/MS spectra was done manually but was assisted by various tools as follows. Manual inspection of the spectrum usually allowed us to derive a tentative sequence. This was used to screen the chlamydial protein sequences NQRA and PqQC (UniProtKB/Swiss-Prot accession numbers O84639 and O84616, respectively) for a possible match using the Mascot software, version 2.2 (Matrix Science Inc, Boston, MA). The MS-product tool (version 5.1.8) (University of California San Francisco), which generates a list of theoretical fragment ions from a given peptide sequence, was then used to match the candidate sequences from the chlamydial proteins to our experimental MS/MS spectra. The assigned sequences were formally confirmed by comparing the experimental MS/MS spectra with those of the corresponding synthetic peptides.

Homology Searches—The sequence homology between chlamydial peptides and human proteins was analyzed in the non-redundant protein database using the Blastp software (National Center for Biotechnology Information, Bethesda, MD). The most recent versions of the database (Release 33, January 22, 2009) and search algorithm available were used.

RESULTS

Identification of Endogenously Processed HLA-B27-restricted T-cell Epitopes from *C. trachomatis* in Human Cells—

The following strategy (Fig. 1) was originally developed in a previous study from our laboratory (16) and was here adapted to investigate the endogenous processing and presentation of HLA-B27 ligands from *C. trachomatis* recognized by $CD8^+$ T-cells from ReA patients (6). 1) B*2705-C1R cells were used as recipients to generate stable transfectants expressing bacterial fusion proteins. 2) To achieve efficient expression of chlamydial proteins in human cells, fusion constructs were generated in which the GFP coding sequence was placed at the 5'-end of the corresponding chlamydial gene. 3) B*2705-bound peptide repertoires were isolated from large amounts of cells expressing or not expressing the bacterial protein. 4) Both peptide pools were fractionated by HPLC in consecutive runs and identical conditions to minimize alterations in the peptide elution patterns. 5) The MALDI-TOF MS spectrum of every HPLC fraction from each peptide pool was obtained and compared with the correlative (i), previous ($i - 1$), and following ($i + 1$) fractions of the other pool to look for peptides found only in the bacterial transfectant. 6) The amino acid sequences of the differential peptides were determined from the corresponding MS/MS spectra. 7) The assigned sequences were validated by comparison with the MS/MS spectra of the corresponding synthetic peptides. If this analysis failed to reveal the presence of the predicted T-cell epitope, a blind MS/MS fragmentation analysis was carried out in the HPLC fraction corresponding to the ex-




FIG. 2. Expression of chlamydial fusion proteins in B*2705-C1R cells. *A*, schematic structure of the GFP-NQRA and GFP-PqqC fusion proteins. *B*, flow cytometry analysis showing the GFP-associated fluorescence of the NQRA (left panel) and PqqC (right panel) fusion protein transfectants. Untransfected B*2705-C1R cells (white histogram) or cells transfected with GFP alone (black histogram) were included as controls. *C*, Western blot analysis showing the expression of the NQRA (left) and PqqC (right) fusion proteins in the respective transfectant cells. The immunoblot was done on the material immunoprecipitated with anti-GFP antibody (NQRA) or on whole lysates (PqqC). In the former case, the prominent nonspecific band corresponds to the heavy chain of the anti-GFP antibody used for immunoprecipitation. In both cases the fusion protein and GFP are noted with upper and lower arrows, respectively.

pected retention time of the peptide epitope and in neighbor fractions using a narrow gate corresponding to the m/z of the peptide.

Expression of Chlamydial Fusion Proteins in B*2705-C1R Cells—We focused on the CT634 and CT610 genes, coding for the NQRA and PqqC proteins, respectively (27), which contain sequences recognized *in vitro* as exogenously added synthetic peptides by CD8⁺ T-cells from ReA patients (6). In addition, PqqC stimulated IFN- γ secretion by peripheral blood mononuclear cells from *Chlamydia*-infected patients with urogenital infection (28). NQRA is a polypeptide of 465 residues involved in Na⁺ transport from the cytoplasm to the periplasm. PqqC is a polypeptide of 231 residues that is expressed in the native state as a homodimer. It is a *Chlamydia*-specific toxin that is secreted into the host cytoplasm and induces apoptosis in mammalian cell lines (29, 30). Both proteins are natural components of elementary bodies (31). GFP-chlamydial gene constructs (Fig. 2A) were transfected into B*2705-C1R cells. The stable expression of the fusion proteins was established by flow cytometry (Fig. 2B) and Western blot analyses (Fig. 2C).

A Single HLA-B27 Chlamydial Ligand Differentially Detected in NQRA Transfectants Is Unrelated to the Predicted T-cell Epitope—HLA-B27-bound peptide pools were isolated from B*2705-C1R cells, B*2705-C1R transfected with GFP alone, and B*2705-C1R expressing the NQRA fusion protein. A total of 4895 ion peaks from the NQRA transfectant were compared with 4958 and 4526 ion peaks from untransfected B*2705-C1R and GFP transfectants, respectively. Only one ion peak from the NQRA transfectant, with m/z 1054.7 in HPLC fr. 208, was absent from B*2705-C1R cells (Fig. 3A) and from the transfectant expressing GFP alone (not shown). The amino acid sequence of this peptide was determined by MALDI-TOF/TOF MS/MS and assigned as a nonamer, KRALLEIVI, spanning residues 86–94 of the chlamydial NQRA protein. The assignment was confirmed by comparison of the MS/MS spectrum with that of the corresponding synthetic peptide (Fig. 3B). The NQRA-(86–94) peptide was unrelated to the predicted T-cell epitope from this protein spanning residues 330–338 (MRDHTITLL) (6). The endogenously processed peptide was not predicted in that report.




FIG. 3. A single endogenous HLA-B27 ligand identified from the chlamydial NQRA protein is unrelated to the predicted T-cell epitope. A, MALDI-TOF MS spectra of fr. 208 from the HLA-B27-bound peptide pools isolated from the NQRA transfectant (above) and untransfected B*2705-C1R cells (below) showing a single ion peak, at m/z 1054.7, differentially present in the bacterial transfectant. Only the relevant sections of the spectra are shown. This ion peak was also not detected in the adjacent fractions 207 and 209 of the untransfected cells (not shown). B, MALDI-TOF/TOF MS/MS spectrum of the ion peak at m/z 1054.7 from the bacterial NQRA transfectant showing the assigned sequence (above) and MS/MS spectrum of the corresponding synthetic peptide (below). The assignment of the ion peak with m/z 129.1 as b₁ is ambiguous because it is also consistent with a Lys immonium ion.

The possibility that low amounts of NQRA-(330–338) might have escaped detection in our comparison was addressed by determining the elution position of the synthetic peptide in the same chromatographic conditions used for fractionation of the HLA-B27-bound peptide pools and looking for the corresponding ion peak in the MALDI-TOF MS spectra of that and neighbor (± 6) HPLC fractions from the bacterial transfectant. No evidence for an ion peak consistent with NQRA-(330–338) was found within the detection limits of the instrument used (data not shown). Of note, this peptide was very inefficient in inducing HLA-B27 refolding *in vitro* (7), suggesting that it is a low affinity ligand.

The Predicted T-cell Epitope from the Chlamydial PqqC Protein Is Endogenously Processed and Presented by HLA-B27—The same strategy was used for determining the endogenous processing of HLA-B27 ligands from PqqC. Upon

systematic comparison of the MALDI-TOF MS spectra from the HPLC-fractionated HLA-B27-bound peptide pools involving 4905 ion peaks from the PqqC transfectant, two ion peaks not found in the corresponding HPLC fractions from B*2705-C1R cells (Figs. 4A and 5A) or B*2705-C1R expressing GFP alone (data not shown) were detected. One of these peaks, with m/z 1055.7 in HPLC fr. 184 from the PqqC transfectant (Fig. 4A), had the same molecular mass and chromatographic retention time as the predicted T-cell epitope from this protein spanning residues 70–78, ARKLLLDNL (6). The amino acid sequence of the differentially expressed B*2705 ligand was determined by MALDI-TOF/TOF MS/MS, and its assignment as the chlamydial T-cell epitope was formally confirmed by comparison of the MS/MS spectrum with that of the corresponding synthetic peptide (Fig. 4B).




FIG. 4. A single endogenous HLA-B27 ligand identified from the chlamydial PqqC protein shows identity with the predicted T-cell epitope. A, MALDI-TOF MS spectra of fr. 184 from the HLA-B27-bound peptide pools isolated from the PqqC transfectant (*above*) and untransfected B*2705-C1R cells (*below*) showing a single detected ion peak, at m/z 1055.7, differentially present in the bacterial transfectant. Only the relevant sections of the spectra are shown. This ion peak was also not detected in the adjacent fractions 183 and 185 of the untransfected cells (not shown). The ion peak at m/z 1053.6, labeled with an asterisk, was a shared ligand found in the adjacent fr. 185 from B*2705-C1R. B, MALDI-TOF/TOF MS/MS spectrum of the ion peak at m/z 1055.7 from the bacterial PqqC transfectant showing the assigned sequence (*above*) and MS/MS spectrum of the corresponding synthetic peptide (*below*). The periodically spaced ion peaks in the *upper* MS/MS spectrum are related to each other by 44 m/z units with a related series differing by 16 m/z units. They arise from traces of the Igepal CA-630 detergent (octylphenyl polyethylene glycol, $(C_2H_4O)_n C_{14}H_{22}O$) used for cell lysis.

The second differential ion peak, with m/z 1183.5, was found in HPLC fr. 182 from the PqqC transfectant (Fig. 5A). The amino acid sequence of this peptide was determined to be TRFSYAEYF (Fig. 5B). It was unrelated to the chlamydial PqqC protein but matched exactly the N-terminal region of the human ankyrin repeat and SOCS box protein 2 (ASB2; UniProtKB/Swiss-Prot accession number Q96Q27). This is a likely substrate recognition component of E3 ubiquitin-protein ligase complexes (32).

NQRA-(86–94) and PqqC-(70–78) Are Highly Homologous to Human Protein Sequences with the HLA-B27 Binding Motif—The chlamydial B*2705 ligand NQRA-(86–94) was screened for homologous sequences against the human proteome database. This search identified a highly homologous sequence (Table I) from the DnaJ homolog DJB14 (UniProtKB/Swiss-Prot accession number Q8TBM8). The follow-

ing features confer special significance to this homology. First, the human counterpart contains the binding motif of HLA-B27, including Arg-2, a nonpolar C-terminal residue (Met-9), and a small N-terminal residue (Ala-1) that is frequent among B*2705 ligands (33). Second, six of nine consecutive residues in the human homolog are identical to the chlamydial peptide. These include Arg-2, the secondary anchor residue Ala-3, and the central region (residues 4–7), which is recognized by the T-cell antigen receptor (TCR) (34). Third, this stretch of identical residues is followed by two conservative differences at positions 8 and 9 involving aliphatic residues.

An analogous search was performed for the chlamydial PqqC-(70–78) epitope. Again this peptide showed a very high homology with residues 172–180 of the exportin 6 (UniProtKB/Swiss-Prot accession number Q96QU8). This human sequence also shows six of nine identical residues, including the




FIG. 5. Identification of a self-derived HLA-B27 ligand differentially detected in PqqC transfectant cells. A, MALDI-TOF MS spectra of fr. 182 from the HLA-B27-bound peptide pools isolated from the PqqC transfectant (*above*) and untransfected B*2705-C1R cells (*below*) showing an ion peak, at m/z 1183.5, differentially present in the bacterial transfectant. Only the relevant sections of the spectra are shown. This ion peak was also not detected in the adjacent fractions 181 and 183 of the untransfected cells (not shown). The ion peaks at m/z 1171.1 and 1205.6 of the bacterial transfectant, labeled with an *asterisk*, were shared ligands found in an independent preparation and in the adjacent fraction 183 from B*2705-C1R, respectively (not shown). B, MALDI-TOF/TOF MS/MS spectrum of the ion peak at m/z 1183.5 from the bacterial PqqC transfectant showing the assigned sequence (*above*) and MS/MS spectrum of the corresponding synthetic peptide (*below*).

TABLE I
Homology of naturally processed HLA-B27 ligands from *C. trachomatis* with human protein sequences

Sequence ^a	Identity %	Organism	Parental protein (residues)	UniProtKB/Swiss-Prot accession no.
KRALLEIVI		<i>C. trachomatis</i>	Na ⁺ -translocating NADH-quinone reductase subunit A, NQRA-(86–94)	O84639
ARALLEIIM	66.7	<i>Homo sapiens</i>	DnaJ homolog protein, DNAJB14-(43–51)	Q8TBM8
ARKLLLDNL		<i>C. trachomatis</i>	Pyroloquinoline-quinone synthase protein, PqqC-(70–78)	O84616
LRKLLLDQV	66.7	<i>H. sapiens</i>	Exportin-6 protein, Exp6-(172–180)	Q96QU8

^a Residues in the human sequences identical to the bacterial peptides are underlined.

Arg-2 motif of HLA-B27 ligands, the secondary anchor residue Lys-3, and identity in the whole central region. The two sequences also show chemically similar aliphatic residues at the anchor positions 1 and 9 and polar residues at the accessible position 8 (Table I). Thus, both endogenously processed HLA-B27 ligands from *C. trachomatis* are highly homologous

to human sequences containing the HLA-B27 binding motif and many features that might favor TCR cross-recognition.

DISCUSSION

The recognition of HLA-B27-restricted chlamydial antigens by CTL is potentially important in the pathogenesis of spon-

dyloarthropathies at least for two reasons. First because it may favor the establishment of bacterial persistence, a crucial event in chronic infection, through the production of IFN- γ (10). Second because it might induce autoimmune cross-reactivity and tissue injury through molecular mimicry with self-derived HLA-B27 ligands (13, 35, 36). Thus, the report of a series of chlamydial HLA-B27 ligands, selected from the screening of the bacterial genome with predictive algorithms, that were recognized by CD8⁺ T-cells from patients with ReA (6, 7) was a significant step toward defining the nature of the CTL responses in this disease. However, this approach has several limitations. The most important one is that, because of the intrinsic T-cell cross-reactivity (21, 22), a synthetic peptide recognized by CTL *in vitro* could be unrelated to the natural epitope that stimulated that CTL *in vivo*. Moreover the predictive algorithms were based on parameters that may not be highly accurate because of insufficient refinement of MHC class I binding motifs (37) or to lack of correspondence between proteasomal cleavage *in vivo* and *in vitro* (38). Finally the screening of the chlamydial genome with these predictive algorithms was limited to nonameric sequences, although a significant percentage of HLA-B27 ligands are longer (33). Thus, the relevance of the predicted T-cell epitopes must be validated by demonstrating their endogenous processing and presentation *in vivo*.

Our approach to this issue involved the expression of chlamydial proteins in human cells and the identification of endogenously processed bacterial HLA-B27 ligands by MS. Several methodological aspects deserve consideration. First, is this a sound alternative to the identification of chlamydial epitopes from infected cells? Second, are the proteins analyzed likely to reach the MHC class I antigen processing pathway under conditions of natural infection? Third, is an MS-based approach sensitive enough to detect natural T-cell epitopes, which might be expressed in very low amounts and still recognized by CTL (39, 40)? The two first questions can only be addressed by considering the mechanism of antigen cross-presentation (41). Chlamydial proteins should reach the MHC class I pathway of the professional antigen-presenting dendritic cells after uptake of antigen-loaded apoptotic bodies. Bacterial proteins in the endosomal compartment of dendritic cells can be transferred to the cytosol by incompletely characterized mechanisms (42, 43) and incorporated into the proteasome pathway. In principle, any bacterial protein might be processed in this way, although many unknown factors in the endosome-to-cytosol transfer mechanism may impose unforeseen limitations on the chlamydial proteins amenable to proteasomal degradation. Other mechanisms that bypass protein entry into the cytosol may also play a role, such as endoplasmic reticulum-phagosome fusion or antigen processing in the endocytic compartment followed by peptide interchange by the MHC class I molecule upon recycling between the endosomal compartment and the cell surface. However, the actual contribution of these pathways to cross-

presentation remains ill defined (41). Thus, the generation of HLA-B27 epitopes upon processing of endogenously synthesized chlamydial proteins may be similar to antigen processing through the cytosolic pathway as it occurs on antigen-presenting cells upon natural infection. However, we cannot rule out that processing of a fusion protein in a B cell line, as analyzed in this study, might differ from processing of a natural bacterial protein in dendritic cells. In *Chlamydia*-infected host cells the most obvious candidates to reach the cytosolic pathway for MHC class I-mediated antigen presentation are the proteins secreted into the cytoplasm (44) or those in the vacuolar membrane (18, 19). However, other chlamydial proteins or protein fragments might reach the cytosol through cross-presentation mechanisms that remain largely undefined in non-professional antigen-presenting cells.

Another question relates to the sensitivity of MS relative to T-cell recognition. We grossly estimated that, in our experimental conditions, the detection limit for HLA-B27 ligands would be around 100–200 copies per cell. This might be 1–2 orders of magnitude below the sensitivity of CTL, which might recognize only a few epitope copies per cell (39, 40). Yet it is reasonable to assume that a transfected protein will yield much more peptide copies presented by HLA-B27 than on infected cells because a much greater protein amount will reach the cytosolic degradation pathway. Thus, it is likely that our system can detect peptides that might be presented upon natural infection in extremely low amounts. For this reason, a candidate chlamydial peptide not found in the B27-bound peptide pool in our analysis is unlikely to be presented on infected cells. Yet this possibility cannot be ruled out.

With the above considerations in mind, our results demonstrate both the reach and the limitations of the epitope prediction algorithms used on the chlamydial genome (6). Although they identified at least one peptide, PqqC-(70–78), they failed to predict another, NQRA-(86–94), and predicted an epitope that is apparently not processed *in vivo*, NQRA-(330–338). Thus, predictive algorithms for putative T-cell epitopes can only be considered as a first step to orient an experimental search for naturally processed antigens. To our knowledge, PqqC-(70–78) is the first HLA-B27-restricted antigen from *C. trachomatis* recognized by CD8⁺ T-cells from ReA patients that has been shown to be endogenously processed and presented *in vivo*. Yet this does not definitely prove that PqqC-(70–78) is a target of the immune response to *C. trachomatis*, which would require showing that the same peptide is presented on *Chlamydia*-infected cells or that a peptide-specific T-cell clone recognizes such cells.

We have previously shown that a peptide derived from the chlamydial DNA primase with high homology to a natural self-derived ligand of HLA-B27 is endogenously processed and presented by this molecule, providing a sound basis for molecular mimicry between chlamydial and self-derived HLA-

B27 ligands (16). The two chlamydial peptides endogenously processed and presented by HLA-B27 in the present study have closely related human sequences, including the same or similar anchor residues and identical TCR-interacting regions. This level of homology between short bacterial peptides and human proteins may be relatively frequent and not restricted to these examples (45). Whether the human homologs are actually processed as natural HLA-B27 ligands is a question of obvious relevance that was not addressed in this study. In the absence of a complete examination of this issue the significance of these homologies is unknown. However, they show that there are multiple possibilities for the generation of self-derived HLA-B27 ligands with obvious mimicry of naturally processed chlamydial epitopes.

We do not know the reason for the differential expression of a self-derived peptide in the PqqC transfectant. There are three possibilities. First, the peptide could be a shared ligand that was detected by chance only in this transfectant but not in the other cells analyzed. Second, the parental ASB2 protein of this self-derived ligand was up-regulated by chance in the particular PqqC transfectant used in our study. Third, up-regulation of ASB2 might have occurred as a specific consequence of the expression of the GFP-PqqC fusion protein. We consider the first possibility unlikely because the differential ion peak was observed in the same transfectant but not in any of the others in two distinct experiments involving independently obtained peptide pools. We cannot rule out the second possibility as it would require repeating this analysis in a second, independently obtained PqqC transfectant, which was not attempted. The third possibility would be consistent with the speculation that GFP-PqqC expression might up-regulate certain components of the ubiquitin/proteasome pathway of protein degradation to which ASB2 belongs. Because this issue is unrelated to the main point of this study, it was not further pursued.

In conclusion, expression of chlamydial proteins in HLA-B27-positive human cells is a reliable and sensitive way to identify endogenously processed bacterial ligands presented by HLA-B27 *in vivo*. Using this approach we demonstrated, for the first time to our knowledge, the endogenous processing and presentation of an HLA-B27-restricted epitope recognized by CD8⁺ T-cells from ReA patients but failed to confirm the endogenous generation of a second T-cell epitope identified by predictive algorithms. In contrast, we identified a non-predicted nonamer from the same parental protein that was endogenously processed and presented by HLA-B27, revealing the limitations of predictive methods for identifying natural MHC class I ligands and T-cell epitopes. The very close similarity of the two bacterial HLA-B27 ligands identified in this study to amino acid sequences containing the HLA-B27 binding motif in the human proteome suggests that this is a possible source of self-derived peptides mimicking chlamydial epitopes. This would be consistent with an autoimmune mechanism of ReA in which T-cell cross-reactivity

between *Chlamydia*-specific and self-derived HLA-B27 ligands may play a crucial role.

Acknowledgment—We thank the staff of the proteomics facilities at the Centro Nacional de Biotecnología, Madrid (member of the ProteoRed network) for help in MS.

* This work was supported by Ministry of Science and Innovation Grants SAF2005/03188 and SAF2008/00461 and an institutional grant from the Fundación Ramón Areces (to the Centro de Biología Molecular Severo Ochoa).

§ The on-line version of this article (available at <http://www.mcponline.org>) contains supplemental material.

‡ To whom correspondence should be addressed: Centro de Biología Molecular Severo Ochoa, c/ Nicolás Cabrera N. 1, Universidad Autónoma, 28049 Madrid, Spain. Tel.: 34-91-196-4554; Fax: 34-91-196-4420; E-mail: aldecastro@cbm.uam.es.

REFERENCES

1. Zeidler, H., Kuipers, J., and Köhler, L. (2004) Chlamydia-induced arthritis. *Curr. Opin. Rheumatol.* **16**, 380–392
2. Zhong, G., Fan, T., and Liu, L. (1999) Chlamydia inhibits interferon gamma-inducible major histocompatibility complex class II expression by degradation of upstream stimulatory factor 1. *J. Exp. Med.* **189**, 1931–1938
3. Zhong, G., Liu, L., Fan, T., Fan, P., and Ji, H. (2000) Degradation of transcription factor RFX5 during the inhibition of both constitutive and interferon gamma-inducible major histocompatibility complex class I expression in chlamydia-infected cells. *J. Exp. Med.* **191**, 1525–1534
4. Zhong, G., Fan, P., Ji, H., Dong, F., and Huang, Y. (2001) Identification of a chlamydial protease-like activity factor responsible for the degradation of host transcription factors. *J. Exp. Med.* **193**, 935–942
5. Loomis, W. P., and Starnbach, M. N. (2002) T cell responses to Chlamydia trachomatis. *Curr. Opin. Microbiol.* **5**, 87–91
6. Kuon, W., Holzhütter, H. G., Appel, H., Grolms, M., Kollnberger, S., Traeder, A., Henklein, P., Weiss, E., Thiel, A., Lauster, R., Bowness, P., Radbruch, A., Kloetzel, P. M., and Sieper, J. (2001) Identification of HLA-B27-restricted peptides from the *Chlamydia trachomatis* proteome with possible relevance to HLA-B27-associated diseases. *J. Immunol.* **167**, 4738–4746
7. Appel, H., Kuon, W., Kuhne, M., Wu, P., Kuhlmann, S., Kollnberger, S., Thiel, A., Bowness, P., and Sieper, J. (2004) Use of HLA-B27 tetramers to identify low-frequency antigen-specific T cells in Chlamydia-triggered reactive arthritis. *Arthritis Res. Ther.* **6**, R521–534
8. Lampe, M. F., Wilson, C. B., Bevan, M. J., and Starnbach, M. N. (1998) Gamma interferon production by cytotoxic T lymphocytes is required for resolution of Chlamydia trachomatis infection. *Infect. Immun.* **66**, 5457–5461
9. Shemer, Y., and Sarov, I. (1985) Inhibition of growth of Chlamydia trachomatis by human gamma interferon. *Infect. Immun.* **48**, 592–596
10. Brunham, R. C., and Rey-Ladino, J. (2005) Immunology of Chlamydia infection: implications for a Chlamydia trachomatis vaccine. *Nat. Rev. Immunol.* **5**, 149–161
11. Keat, A. C., Maini, R. N., Nkwazi, G. C., Pegrum, G. D., Ridgway, G. L., and Scott, J. T. (1978) Role of Chlamydia trachomatis and HLA-B27 in sexually acquired reactive arthritis. *Br. Med. J.* **1**, 605–607
12. Young, J. L., Smith, L., Matyszak, M. K., and Gaston, J. S. (2001) HLA-B27 expression does not modulate intracellular Chlamydia trachomatis infection of cell lines. *Infect. Immun.* **69**, 6670–6675
13. Bachmaier, K., Neu, N., de la Maza, L. M., Pal, S., Hessel, A., and Penninger, J. M. (1999) Chlamydia infections and heart disease linked through antigenic mimicry. *Science* **283**, 1335–1339
14. Popov, I., Dela Cruz, C. S., Barber, B. H., Chiu, B., and Inman, R. D. (2002) Breakdown of CTL tolerance to self HLA-B*2705 induced by exposure to Chlamydia trachomatis. *J. Immunol.* **169**, 4033–4038
15. Ramos, M., Alvarez, I., Sesma, L., Logean, A., Rognan, D., and López de Castro, J. A. (2002) Molecular mimicry of an HLA-B27-derived ligand of arthritis-linked subtypes with chlamydial proteins. *J. Biol. Chem.* **277**, 37573–37581
16. Cragolini, J. J., and de Castro, J. A. (2008) Identification of endogenously

- presented peptides from *Chlamydia trachomatis* with high homology to human proteins and to a natural self-ligand of HLA-B27. *Mol. Cell. Proteomics* **7**, 170–180
17. Karunakaran, K. P., Rey-Ladino, J., Stoykov, N., Berg, K., Shen, C., Jiang, X., Gabel, B. R., Yu, H., Foster, L. J., and Brunham, R. C. (2008) Immunoproteomic discovery of novel T cell antigens from the obligate intracellular pathogen *Chlamydia*. *J. Immunol.* **180**, 2459–2465
 18. Fling, S. P., Sutherland, R. A., Steele, L. N., Hess, B., D'Orazio, S. E., Maisonneuve, J., Lampe, M. F., Probst, P., and Starnbach, M. N. (2001) CD8+ T cells recognize an inclusion membrane-associated protein from the vacuolar pathogen *Chlamydia trachomatis*. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 1160–1165
 19. Starnbach, M. N., Loomis, W. P., Owendale, P., Regan, D., Hess, B., Alderson, M. R., and Fling, S. P. (2003) An inclusion membrane protein from *Chlamydia trachomatis* enters the MHC class I pathway and stimulates a CD8+ T cell response. *J. Immunol.* **171**, 4742–4749
 20. Grotenbreg, G. M., Roan, N. R., Guillen, E., Meijers, R., Wang, J. H., Bell, G. W., Starnbach, M. N., and Ploegh, H. L. (2008) Discovery of CD8+ T cell epitopes in *Chlamydia trachomatis* infection through use of caged class I MHC tetramers. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 3831–3836
 21. Wucherpfennig, K. W., and Strominger, J. L. (1995) Molecular mimicry in T cell-mediated autoimmunity: viral peptides activate human T cell clones specific for myelin basic protein. *Cell* **80**, 695–705
 22. Evavold, B. D., Sloan-Lancaster, J., Wilson, K. J., Rothbard, J. B., and Allen, P. M. (1995) Specific T cell recognition of minimally homologous peptides: evidence for multiple endogenous ligands. *Immunity* **2**, 655–663
 23. Zemmour, J., Little, A. M., Schendel, D. J., and Parham, P. (1992) The HLA-A,B “negative” mutant cell line C1R expresses a novel HLA-B35 allele, which also has a point mutation in the translation initiation codon. *J. Immunol.* **148**, 1941–1948
 24. Paradela, A., García-Peydró, M., Vázquez, J., Rognan, D., and López de Castro, J. A. (1998) The same natural ligand is involved in all recognition of multiple HLA-B27 subtypes by a single T cell clone: role of peptide and the MHC molecule in alloreactivity. *J. Immunol.* **161**, 5481–5490
 25. Barnstable, C. J., Bodmer, W. F., Brown, G., Galfre, G., Milstein, C., Williams, A. F., and Ziegler, A. (1978) Production of monoclonal antibodies to group A erythrocytes, HLA and other human cell surface antigens. New tools for genetic analysis. *Cell* **14**, 9–20
 26. Paradela, A., Alvarez, I., García-Peydró, M., Sesma, L., Ramos, M., Vázquez, J., and López De Castro, J. A. (2000) Limited diversity of peptides related to an alloreactive T cell epitope in the HLA-B27-bound peptide repertoire results from restrictions at multiple steps along the processing-loading pathway. *J. Immunol.* **164**, 329–337
 27. Stephens, R. S., Kalman, S., Lammel, C., Fan, J., Marathe, R., Aravind, L., Mitchell, W., Olinger, L., Tatusov, R. L., Zhao, Q., Koonin, E. V., and Davis, R. W. (1998) Genome sequence of an obligate intracellular pathogen of humans: *Chlamydia trachomatis*. *Science* **282**, 754–759
 28. Olsen, A. W., Follmann, F., Højrup, P., Leah, R., Sand, C., Andersen, P., and Theisen, M. (2007) Identification of human T cell targets recognized during *Chlamydia trachomatis* genital infection. *J. Infect. Dis.* **196**, 1546–1552
 29. Stenner-Liewen, F., Liewen, H., Zapata, J. M., Pawlowski, K., Godzik, A., and Reed, J. C. (2002) CADD, a *Chlamydia* protein that interacts with death receptors. *J. Biol. Chem.* **277**, 9633–9636
 30. Schwarzenbacher, R., Stenner-Liewen, F., Liewen, H., Robinson, H., Yuan, H., Bossy-Wetzel, E., Reed, J. C., and Liddington, R. C. (2004) Structure of the *Chlamydia* protein CADD reveals a redox enzyme that modulates host cell apoptosis. *J. Biol. Chem.* **279**, 29320–29324
 31. Skipp, P., Robinson, J., O'Connor, C. D., and Clarke, I. N. (2005) Shotgun proteomic analysis of *Chlamydia trachomatis*. *Proteomics* **5**, 1558–1573
 32. Heuzé, M. L., Guibal, F. C., Banks, C. A., Conaway, J. W., Conaway, R. C., Cayre, Y. E., Benecke, A., and Lutz, P. G. (2005) ASB2 is an Elongin BC-interacting protein that can assemble with Cullin 5 and Rbx1 to reconstitute an E3 ubiquitin ligase complex. *J. Biol. Chem.* **280**, 5468–5474
 33. Lopez de Castro, J. A., Alvarez, I., Marcilla, M., Paradela, A., Ramos, M., Sesma, L., and Vázquez, M. (2004) HLA-B27: a registry of constitutive peptide ligands. *Tissue Antigens* **63**, 424–445
 34. Garboczi, D. N., Ghosh, P., Utz, U., Fan, Q. R., Biddison, W. E., and Wiley, D. C. (1996) Structure of the complex between human T-cell receptor, viral peptide and HLA-A2. *Nature* **384**, 134–141
 35. Benjamin, R., and Parham, P. (1990) Guilt by association: HLA-B27 and ankylosing spondylitis. *Immunol. Today* **11**, 137–142
 36. Bachmaier, K., and Penninger, J. M. (2005) *Chlamydia* and antigenic mimicry. *Curr. Top. Microbiol. Immunol.* **296**, 153–163
 37. Ruppert, J., Sidney, J., Celis, E., Kubo, R. T., Grey, H. M., and Sette, A. (1993) Prominent role of secondary anchor residues in peptide binding to HLA-A2.1 molecules. *Cell* **74**, 929–937
 38. Rock, K. L., and Goldberg, A. L. (1999) Degradation of cell proteins and the generation of MHC class I-presented peptides. *Annu. Rev. Immunol.* **17**, 739–779
 39. Sykulev, Y., Joo, M., Vturina, I., Tsomides, T. J., and Eisen, H. N. (1996) Evidence that a single peptide-MHC complex on a target cell can elicit a cytolytic T cell response. *Immunity* **4**, 565–571
 40. Purbhoo, M. A., Irvine, D. J., Huppa, J. B., and Davis, M. M. (2004) T cell killing does not require the formation of a stable mature immunological synapse. *Nat. Immunol.* **5**, 524–530
 41. Lin, M. L., Zhan, Y., Villadangos, J. A., and Lew, A. M. (2008) The cell biology of cross-presentation and the role of dendritic cell subsets. *Immunol. Cell Biol.* **86**, 353–362
 42. Ackerman, A. L., Giardini, A., and Cresswell, P. (2006) A role for the endoplasmic reticulum protein retrotranslocation machinery during cross-presentation by dendritic cells. *Immunity* **25**, 607–617
 43. Rock, K. L. (2006) Exiting the outside world for cross-presentation. *Immunity* **25**, 523–525
 44. Kleba, B., and Stephens, R. S. (2008) Chlamydial effector proteins localized to the host cell cytoplasmic compartment. *Infect. Immun.* **76**, 4842–4850
 45. Davies, J. M. (2000) Introduction: Epitope mimicry as a component cause of autoimmune disease. *Cell Mol. Life Sci.* **57**, 523–526