

File: Membrane Proteome
Date: October 6, 2010

Comprehensive Proteomic Analysis of Membrane Proteins in *Toxoplasma gondii*

Fa-Yun Che^{1,2}; Carlos Madrid-Aliste^{2,3}; Berta Burd²; Hongshan Zhang²; Edward Nieves^{2,5}; Kami Kim^{2,6,7}; Andras Fiser^{2,3,5}; Ruth Hogue Angeletti^{2,4,5}; Louis M. Weiss^{1,2,6}

¹ Department of Pathology,

² Biodefense Proteomics Research Center,

³ Department of Systems and Computational Biology,

⁴ Department of Developmental and Molecular Biology,

⁵ Department of Biochemistry

⁶ Department of Medicine (Infectious Diseases),

⁷ Department of Microbiology and Immunology,

Albert Einstein College of Medicine, Bronx, NY 10461

§ Corresponding authors: Fa-Yun Che and Louis M. Weiss

Albert Einstein College of Medicine
1300 Morris Park Ave. Bronx, NY 10461.

Phone: 718 430 2142

Fax: 718 430 8543

E-mail: fa-yun.che@einstein.yu.edu, louis.weiss@einstein.yu.edu

Running Title: Membrane proteome of *T. gondii*

Keywords: *Toxoplasma gondii*, proteomics, membrane proteins, three-layer sandwich gel electrophoresis (TLSGE), multidimensional protein identification technology (MudPIT).

SUMMARY

Toxoplasma gondii (*T. gondii*) is an obligate intracellular protozoan parasite that is important human and animal pathogen. Experimental information on *T. gondii* membrane proteins is limited, and the majority of gene predictions with predicted transmembrane motifs are of unknown function. A systematic analysis of the membrane proteome of *T. gondii* is important not only for understanding this parasite's invasion mechanism(s), but also for the discovery of potential drug targets and new preventative and therapeutic strategies. Here we report a comprehensive analysis of the membrane proteome of *T. gondii*, employing three proteomics strategies: 1D gel LC-MS/MS analysis (one-dimensional gel electrophoresis liquid chromatography-tandem mass spectrometry), biotin labeling in conjunction with 1D gel LC-MS/MS analysis, and a novel strategy that combines three-layer “Sandwich” Gel Electrophoresis (TLSGE) with multidimensional protein identification technology (MudPIT). A total of 2,241 *T. gondii* proteins with at least one predicted transmembrane segment were identified and grouped into 841 sequentially non-redundant protein clusters, which account for 21.8% of the predicted transmembrane protein clusters in the *T. gondii* genome. A large portion (42%) of the identified *T. gondii* membrane proteins are hypothetical proteins. Furthermore, many of the membrane proteins validated by mass spectrometry are unique to *T. gondii* or to the Apicomplexa, providing a set of gene predictions ripe for experimental investigation, and potentially suitable targets for the development of therapeutic strategies.

INTRODUCTION

Toxoplasma gondii (*T. gondii*), an obligate intracellular parasitic protozoan, is a common causative agent of disease in immune competent and immune compromised humans (1, 2). It is capable of causing severe congenital neurological impairment if acquired *in utero* (1, 2). Infection can reactivate from latent tissue cysts in patients with immune suppression (AIDS or chemotherapy) resulting in acute infections most often manifesting as encephalitis. Toxoplasmosis is a major cause of death among patients with AIDS (3, 4). The development of new preventative and therapeutic strategies relies on an improved understanding of the interaction between pathogens and their hosts (5). Reported *T. gondii* infection rates can be as high as 70%, depending on the populations or geographic area studied. *T. gondii* can infect all warm-blooded mammals although the definitive hosts are members of the cat family.

There are two phases in the parasite's life cycle, the sexual phase, which takes place only in felines, and the asexual phase, which takes place in any warm-blooded animal. *T. gondii* has three functionally distinct pathogenic forms, sporozoites (in oocysts), tachyzoites and bradyzoites (in tissue cysts) (6). When tissue cysts (*e.g.*, from an infected mouse) are ingested by a cat, the cysts survive the passage through the stomach to infect the epithelial cells of the small intestine, where they differentiate and reproduce sexually eventually forming oocysts, which are then shed with the feces. The oocysts contain sporozoites that develop into tachyzoites upon ingestion by mammals. In a similar fashion, if tissue cysts (which contain bradyzoites) are ingested by other mammals they can differentiate into tachyzoites. Tachyzoites, the invasive form of *T. gondii*, rapidly multiply asexually, invade host cells, and are distributed via the blood stream and lymphatic system throughout the body including the brain. Tachyzoites are responsible for acute infection

and disease. Tachyzoites differentiate in response to stress, such as the inflammatory response, into bradyzoites, which are contained within tissue cysts. Humans, as intermediate hosts, can be exposed to and infected with *T. gondii* via food-borne, water-borne or maternofetal routes. Maternofetal transmission causes congenital infection which can result in miscarriage, mental retardation, learning disabilities, blindness, microcephaly and seizures, or death (3, 7, 8, 9). Upon invasion of host cells *T. gondii* can commandeer host functions affecting the secretion of cytokines (2) and other cellular processes ultimately bypassing host cell defenses (10, 11). The invasion process involves several steps including the recognition of surface receptors by surface antigens and the secretion of proteins from specialized secretory organelles, such as rhoptries, micronemes and dense granules. Invasion results in the formation of the parasitophorous vacuole membrane (PVM) by invaginating host plasma membrane and selective exclusion of many host integral membrane proteins (10, 11). The PVM envelops and shelters the parasite inside the parasitophorous vacuole and prevents acidification of this compartment, in which parasites can grow and replicate (12). Parasite membrane proteins play critical roles in invasion and interaction with host cells. For example, *T. gondii* apical membrane antigen-1 (TgAMA1), a type I transmembrane protein, localizes to the parasite's micronemes, secretory organelles. TgAMA1 depletion inhibits secretion of the rhoptries whose discharge is coupled to active host cell penetration (13). Besides cell-cell interactions, other functions of membrane proteins include ion and solute transport (14), cell signaling, and catalysis. It is likely that, as in other eukaryotes, targets of successful therapeutic drugs will be identified among these parasite membrane proteins. In addition, analysis of the membrane proteome of *T. gondii* should further improve our understanding of the molecular mechanisms of parasite invasion and pathogenesis and assist in identifying new chemotherapy targets. Several thousand proteins are predicted to be membrane

bound in the *T. gondii* genome according to transmembrane segment prediction algorithms.

Presently, however, only a limited number of membrane proteins have been reported and characterized in this parasite (15, 16), reflecting the challenges in studying membrane proteins due to their high hydrophobicity and low abundance.

Membrane proteins are insoluble in aqueous solution, putting researchers at a dilemma when handling membrane protein samples and carrying out downstream proteomics. Solubilization of membrane proteins in aqueous solution requires the use of detergents (such as SDS or Triton X-100), non-polar solvents, or sometimes denaturing agents. The presence of relatively high concentrations of detergents or denaturing agents in a sample generally results in inefficient enzymatic digestion (due to enzyme inactivation) and reduces the downstream performance of peptide separation on reverse phase chromatography. Furthermore, a major disadvantage of the presence of detergent in the sample is the suppression of peptide ionization and detrimental interference during electrospray ionization. Background ions raised from detergents can obscure the signal in mass spectra. In addition, detergents can cause the formation of adducts and a shift of the charge envelope (17), making the interpretation of mass spectra more difficult. Researchers have developed several variants of acid cleavable detergents, such as PPS (sodium 3-(4-(1, 1-bis(hexyloxy)ethyl)pyridinium-1-yl) propane-1-sulfonate), ProteaseMAX and RapiGest SF, which can be used to help solubilize and unfold membrane proteins to improve enzymatic digestion. After digestion, the detergent undergoes hydrolysis under acidic conditions facilitating proteomic studies, and then is removed from the sample solution.

To overcome the obstacles from using detergents to study membrane proteins, one-dimensional gel electrophoresis coupled liquid chromatography-tandem mass spectrometry analysis (1D gel LC-MS/MS) has proven to be a relatively simple and effective approach (18, 19).

The high concentration of detergents can be removed after membrane proteins are resolved on SDS PAGE gel.

Another approach to membrane protein analysis that specifically targets plasma membrane proteins is biotinylation of cell surface proteins and affinity purification. Cells are labeled with sulfo-NHS-SS-biotin and the labeled proteins are collected with streptavidin beads, resolved on SDS PAGE gel, followed by the typical protein identification procedure using LC-MS/MS (20, 21). Recently, Zhu and colleagues developed a “Tube-Gel” digestion protocol for membrane proteins (22). Membrane proteins in the presence of high concentrations of detergents are incorporated into a polyacrylamide gel matrix prior to casting the gel using a miniaturized tube. The tube gel is removed without electrophoresis and washed to remove detergents before in-gel digestion. The Tube-Gel digestion protocol can be used to solubilize membrane proteins with minimal interference in downstream LC-MS/MS analysis (22). A drawback, however, with Tube-Gel protocol is that acrylamide can chemically modify some amino acid residues in proteins. A novel gel electrophoresis system, Three-layer “Sandwich” Gel Electrophoresis (TLSGE), was recently introduced by Lynn and colleagues for efficient salt removal and concentrating protein (23). A powerful feature of TLSGE is its ability to handle relatively large volumes of sample, even several mL, concentrating proteins on a small piece of protein gel and removing detergents and salts during electrophoresis. It is a very useful enrichment strategy for low abundance membrane proteins. In this paper we report the benefits of combining TLSGE with MudPIT (TLSGE MudPIT) and RapiGest SF for the proteomic analysis of membrane proteins in *T. gondii*.

To comprehensively decipher the membrane proteome of *T. gondii*, we implemented three approaches: 1D gel LC-MS/MS, TLSGE MudPIT, and biotin-directed affinity purification (BDAP) (20, 21). The MS/MS data were analyzed in the context of a variety of alternative gene predictions

from *T. gondii*, and 841 sequentially non-redundant *T. gondii* membrane protein clusters were validated from thousands of predicted membrane proteins. Approximately, 42% of these identified membrane proteins are hypothetical proteins, 50% of which are unique to *T. gondii*.

EXPERIMENTAL PROCEDURES

Reagents

Thioglucopyranoside, Percoll, urea, thiourea, octyl- β -glucoside, dithiothreitol (DTT), iodoacetamide, TCEP (tris(2-carboxyethyl)phosphine), ammonia bicarbonate, Coomassie Brilliant Blue R-250, KCl, KH₂PO₄, H₃PO₄ and Na₂CO₃ were purchased from SIGMA (St. Louis, MO USA). Acetonitrile was from Fisher Scientific (Pittsburgh, PA, USA). CompleteTM Proteinase inhibitor cocktail was purchased from Santa Cruz Biotechnology (Santa Cruz, CA., USA). Trifluoroacetic acid (TFA), formic acid and EZ-LinkTM sulfo-NHS-SS-biotin were purchased from Pierce (Rockford, IL, USA). Sequence grade trypsin was purchased from Promega (Madison, WI, USA). Streptavidin magnetic beads were purchased from BioLabs (Ipswich, MA, USA). RapiGestTMSF was purchased from Waters (Milford, MA, USA). Dulbecco's modified eagle medium (DMEM), Phosphate Buffered Saline (PBS), fetal calf serum, glutamine and penicillin streptomycin were purchased from Invitrogen (Carlsbad, CA, USA).

Growth of Toxoplasma gondii in vitro and Isolation of Tachyzoites

RH strain *T. gondii* parasites were maintained by serial passage in confluent monolayers of human foreskin fibroblasts (HFF) in Dulbecco's modified eagle medium (DMEM) supplemented with 10% fetal calf serum (FBS), 2 mM glutamine and 5 mM penicillin-streptomycin in a 5% CO₂ incubator at 37 °C. The absence of mycoplasma contamination was monitored monthly using a PCR method (Invitrogen) and only mycoplasma negative cells were used for culture. After 4 to 5 days in culture, tachyzoites were counted in a Neubauer hemocytometer, washed three times with PBS and the freshly lysed parasites were scraped off the culture flasks. *T. gondii* was isolated from HFF by passage through with a 25-gauge needle followed by filtration through a Nucleopore filter (3µm pore size) as previously published (24). Parasites were then pelleted by centrifugation.

Purification of Membrane Proteins from Toxoplasma gondii

About 4.0×10^9 purified *T. gondii* RH strain tachyzoites from culture were resuspended in 20 ml of SMDI buffer (250 mM Sucrose, 10 mM MOPS-KOH, pH 7.2, 2 mM DTT, 1× protease inhibitor cocktail) and disrupted by a French press at a pressure of 1000 Psi, medium setting. The lysate was centrifuged at $756 \times g$ at 4°C for 10 min to pellet unbroken cells. Intact parasites and large debris were resuspended in 10 mL of SMDI buffer and disrupted once more by French press at a pressure of 1000 Psi, medium setting. The pooled supernatant was centrifuged at $25,000 \times g$ at 4°C for 20 min. The supernatant was saved for analysis as the cytosolic fraction. The pellet was resuspended in 10 mL of 30 % Percoll in SMDI buffer. After centrifugation at $75,000 \times g$ in an ultracentrifuge (Rotor TLA 100.3; 30,000 rpm) at 4°C for 25 min, the top band was collected from the self-generated gradient. The band was diluted in SMDI buffer and spun at $100,000 \times g$ at 4°C for 90 min (Rotor TLA 100.3; 40,000 rpm). A band collected between the buffer and resultant

Percoll cushion contained the *T. gondii* ghost fraction consisting of membranes and cytoskeleton (24). To isolate the membrane fraction, *T. gondii* ghosts were resuspended in an equal volume of 2 % thioglucoyanoside in 40 mM Tris, pH 7.6, by pipeting the mixture up and down 10 times (suspension kept on ice) followed by a brief vortex. After centrifugation at $20,000 \times g$ in an Eppendorff centrifuge at 4°C for 20 min, the supernatant was saved as the membrane fraction (extraction 1). This extraction was repeated twice with 300 μL of 1% thioglucoyanoside (extraction 2 and 3, respectively). These fractions were pooled and frozen in liquid nitrogen and stored at -80°C until protein analysis. In the parallel experiment, prior to being resuspended in thioglucoyanoside solution, *T. gondii* ghosts were first washed sequentially with 1M KCl and 0.1 M Na_2CO_3 twice.

1D SDS PAGE and In-gel Digestion

For SDS PAGE, approximately 60 μg of proteins was mixed with 50 μL of 2 \times SDS loading buffer and heated at 85°C for 10 min prior to loading on a precast gel (Invitrogen, Carlsbad, CA, USA). Proteins were afterward stained with Coomassie Brilliant Blue R-250. Usually, 40-45 gel bands were cut across the whole sample lane. Each band was excised further into small pieces ($\sim 1 \times 2$ mm) and placed into a 0.65 mL micro-centrifuge siliconized tube (PGC Scientific) and washed with 250 μL of Millipore water twice for 5 min. In-gel trypsin digestion was performed. Gel pieces were washed three times each with an appropriate volume of 25 mM NH_4HCO_3 in 50% acetonitrile. The gel pieces were dried completely in a Speed Vac. The reduction reaction was allowed to proceed at 56°C for 1 h in an appropriate volume of 10 mM DTT in 25 mM NH_4HCO_3 and the alkylation was allowed to proceed in the dark for 45 min at room temperature in an

appropriate volume of 55 mM iodoacetamide in 25 mM NH_4HCO_3 . The gel pieces were washed with 100 μL of 25 mM NH_4HCO_3 for 10 min and dehydrated with 100 μL of 25 mM NH_4HCO_3 in 50% acetonitrile for 5 min. The wash and dehydration step was repeated once. Following drying in a SpeedVac, the gel pieces were mixed with an appropriate volume of 12.5 ng/ μL of trypsin and incubated on ice for 40 min and 25 mM NH_4HCO_3 was added as needed to cover the gel pieces. Digestion was then carried out at 37°C overnight. To extract the tryptic peptides from the gel pieces, an appropriate volume of 60% acetonitrile, 0.2% TFA, was added. Following 20 min of vortex and 5 min of sonication, the supernatant was taken and saved. The extraction was repeated once and all the supernatants were combined (in-gel digestion protocol obtained from <http://ms-facility.ucsf.edu/ingel.html>). After the evaporation of acetonitrile in a SpeedVac, the sample was desalted with a C18 ZipTip (Millipore), and half of the eluate was analyzed with nanoLC-MS/MS.

Preparation of Integral Plasma Membrane Proteins Using Biotin Labeling

T. gondii RH strain tachyzoites ($\sim 4.0 \times 10^9$ cells) collected from cell culture were washed five times in 20 mL PBS (pre-warmed at 37 °C) and resuspended in 20 mL PBS. About 25 mg EZ-LinkTM sulfo-NHS-SS-biotin was added to the mixture and incubated for 30 min at room temperature. The mixture was centrifuged for 20 min at 4150 rpm and 15 °C. The supernatant was removed, and the tachyzoites resuspended in 20 mL PBS containing 1 mg/mL lysine and incubated for 15 min at room temperature. The mixture was centrifuged for 20 min at 4150 rpm at 15 °C and, the supernatant removed. *T. gondii* tachyzoites were washed twice with 20 mL of PBS. The *T. gondii* ghost fraction was prepared as described (see above “Purification of Membrane Proteins

from *Toxoplasma gondii*”). The ghost fraction was resuspended in 500 μ L of 1% thioglucopyranoside in 40 mM Tris, pH 7.6, 1 \times protease inhibitor, containing around 9-10 mg prewashed streptavidin magnetic beads. After 2 h incubation at 4 $^{\circ}$ C, magnetic beads were collected using a magnetic rack (BioLabs). The beads were sequentially washed three times with 1.2 mL of ice-cold 1 M KCl, three times with 1.2 mL of 0.1 M Na_2CO_3 , then once with 1.2 mL of 1% thioglucopyranoside in 40 mM Tris, pH 7.6, 1 \times protease inhibitor. Finally, the beads were suspended in 85 μ L of 2 \times SDS PAGE sample buffer containing 100 mM of DTT, 30 mM biotin, 6 M urea, and 2 M thiourea. Incubation was carried out for 10 min at room temperature followed by 10 min at 95 $^{\circ}$ C. The supernatant was collected after centrifugation for 3 min at 11,000 g. Aliquots of the supernatant were separated with SDS PAGE. 41 gel bands were cut across the whole sample lane. In-gel digestion was performed for each band as described above.

Three-Layer “Sandwich” Gel Electrophoresis (TLSGE), In-gel Digestion and Strong Cation Exchange Chromatography of Tryptic Peptides

The preparation of the three-layer “sandwich” gel (TLSG) was performed as described previously (23). The three-layer “sandwich” gel electrophoresis (TLSGE) was carried out with an electro-eluter device (Bio-Rad, model 422). The bottom layer or sealing layer was first formed by casting 1 mL of 40 % polyacrylamide gel in a glass tube (1cm I.D. \times 6 cm long, Bio-Rad) (Fig. 1). Immediately on top of the sealing layer is cast the concentrating layer with 200~250 μ L of 10 % acrylamide gel. About 500 μ L of 2 \times SDS PAGE sample loading buffer was mixed with 500~1000 μ L of membrane protein fraction. The resulting solution was then mixed with 0.5-1.0 mL of pre-heated 1% agarose in the elution buffer containing 25 mM Tris base, 192 mM Glycine and

0.2 % SDS, and immediately added on top of the concentrating layer forming the upper layer of the TLSG. After the glass tube cooled down, the tube was placed into the electro-eluter with the upper and lower elution buffers, a 150 constant voltage was applied and the electrophoresis was allowed to run for 4-5 hours. When a potential is applied between the upper and lower layers of TLSG, proteins migrate from the agarose gel into the middle layer and are concentrated in this layer; the bottom layer prevents protein leakage from the middle layer due to the high density acrylamide gel (40%) (Fig.1). Consequently, TLSGE concentrates almost all proteins from a solution into the middle layer gel slice through electrophoresis while salts and detergents are removed from this layer gel. After electrophoresis, the concentrating layer of gel was taken out, excised into small pieces (~1×2 mm), and placed into a 1.5 mL micro-centrifuge siliconized tube (PGC Scientific). The gel pieces were washed with 600 μ L of Millipore water for 10 min, and this wash step was repeated at least four or five times. In-gel tryptic digestion was performed according to the procedure described above with some modifications: prior to the addition of trypsin, 200-300 μ L of 0.1% RapiGestTMSF was added to the dried gel pieces before incubating in a 37°C water bath for 30 min, followed by drying the gel pieces completely in a Speed Vac. The extracted tryptic peptide solution was concentrated in a SpeedVac to 300~400 μ L and mixed with 600 μ L of buffer A used during strong cation exchange chromatography (SCX). The resulting sample was vortexed briefly and centrifuged at 13,000 rpm for 10 min. The supernatant was taken, mixed with 9 mL of SCX buffer A and loaded on a 10 mL sample loop of the AKTA purifier (Amersham Biosciences). The SCX separation column was a Polysulfoethyl A column (2.1 mm i.d. ×100 mm, 5 μ m, 300Å, PolyLC Inc.). Buffer A was 10 mM KH₂PO₄, 25% ACN, pH 2.8; buffer B was 10 mM KH₂PO₄, 700 mM KCl, 25 % ACN, pH 2.8. The gradient was from 0 % to 50% buffer B in 120 min and from 50% to 100% B in 20 min with a flow rate of 0.1 mL/min. One fraction was collected every

minute. Each fraction was concentrated to 20 μL in a SpeedVac, desalted with a C18 ZipTip, and half of the eluate was analyzed by nanoLC-MS/MS.

NanoLC-MS/MS Analysis

Nanoelectrospray LC-MS/MS analysis was performed on a linear ion trap mass spectrometer (LTQ, Thermo, San Jose, CA) interfaced with a TriVersa NanoMate nanoelectrospray ion source (Advion BioSciences, Ithaca, NY). An Ultimate^{Plus} nano-HPLC system with a Famos autosampler (Dionex Corporation, Sunnyvale, CA), was coupled with the TriVersa NanoMate. Peptides, 25 μL , were loaded on a C18 μ -PrecolumnTM Cartridge (5 μm , 100 \AA , 300 μm i.d. \times 5mm) by the autosampler with a 25 μL sample loop at a flow rate of 15 $\mu\text{L}/\text{min}$. Mobile phase A was 2 % acetonitrile and 0.1 % formic acid in water and mobile phase B was 80 % acetonitrile and 0.1 % formic acid in water. The HPLC flow rate used was 250 nL/min. After injection of 25 μL sample, and washing for 20 minutes with 100% mobile phase A, the precolumn was switched in line with the analytical column, C18 PepMap100, 3 μm , 100 \AA , 75 μm i.d. \times 150 mm (Dionex Corporation, Sunnyvale, CA). Then mobile phase B was then increased from 2 to 55 % over 70 minutes, held for 5 minutes, increased to 95 % over 20 minutes and held at 95 % B for 5 minutes. The four most intense ions having a charge state between +2 to +4, determined from an initial survey scan from 300-1800 m/z , were selected for zoom scan and MS/MS. In some nanoLC-MS/MS experiments, the ten most intense ions from each initial MS survey were selected for MS/MS only. MS/MS was performed using an isolation width of 2 m/z ; normalized collision energy of 35 % and a minimum signal intensity of 1000 counts. Dynamic exclusion was enabled, such that once a certain ion is selected twice for MS/MS within 30 sec, this ion is excluded from being selected again for MS/MS

during the next 120 sec.

Data and Bioinformatics Analysis

Protein database search of MS/MS spectra DTA files were created from the raw LC-MS/MS data, and searched with Mascot Daemon software (Version 2.2.2) against a database comprised of the human proteome (~40,000 sequences) and predicted *T. gondii* protein sequence sets (38,184 sequences) organized in our Experimental Proteomics DataBase for *T. gondii* and *Cryptosporidium parvum* EPIC-DB (24). EPIC-DB is a comprehensive collection of experimental and computational data that contains all available protein sequences for *T. gondii* and *C. parvum*, which represents the theoretical proteome of the respective organisms. The available *T. gondii* sequences were compiled from five different gene prediction datasets: TigrScan, TwinScan, GlimmerHMM, toxoDB Release 4.30, ME49 (Release 6.0, <http://ToxoDB.org>), and experimentally characterized sequences obtained from the NCBI protein database. The combination of alternative gene predictions was necessary because there was a surprisingly small overlap between any two alternative sets (24). In addition, MS/MS analysis showed that all gene prediction sets contain experimentally validated unique predictions. A false discovery rate (FDR) for peptide identification was assessed by decoy database searching. The following parameters were used for all searches: trypsin, 2 missed cleavages; variable modifications of carbamidomethylation (Cys), deamidation (Asn and Gln) and oxidation (Met); monoisotopic masses; peptide mass tolerance of 3.5 Da; and product ion mass tolerance of 0.6 Da. Proteins were considered identified having at least two bold red peptides (BR, the most logical assignment of a peptide to a proteins and prevents duplicate homologous proteins to be reported) and having a

peptide ion score cut-off of 49 or greater (corresponding to $p < 0.05$). For those proteins identified by only one BR peptide and having a peptide ion score cut-off of 49 or greater (corresponding to $p < 0.05$), the mass spectrometry raw data was checked manually to meet the following criteria: the difference between the measured and theoretical peptide parent mass divided by theoretical one was within 100 ppm; MS/MS product ions within 0.5 Da of the predicted b and y ions; and 80% of the high intensity product ions matched to either b or y ions; and both b and y ions matched. A BLAST search against other genomes was also performed with the peptide hits to rule out possible protein hits from other contaminating species. A list of all significant *T. gondii* protein hits are provided in Supplementary Tables 1, 2 and 3. All references to subsequent protein hits refer to those of *T. gondii* only, unless specified as human.

Clustering protein sequences, trans-membrane predictions Redundant *T. gondii* protein sequences were clustered using a greedy clustering algorithm (CD-HIT) using a 90% sequence identity threshold (24, 26-28). Transmembrane segments were predicted using Phobius, a combined trans-membrane protein topology and signal peptide predictor (29).

GPI prediction and plotting trans-Membrane domains GPI-anchor proteins were predicted using PredGPI (<http://www.biomedcentral.com/1471-2105/9/392>) (30). 2D topology images of transmembrane proteins were created using TOPO2 software (<http://www.sacs.ucsf.edu/TOPO2/>).

GO annotations and functional annotation Homologous sequences in apicomplexan genomes, human genome and the NCBI NR database were identified by a BLAST search (without

low complexity filtering and e-value of 0.001). The homologues sequences found by BLAST in the three dataset were mapped to the GO database by their GI number. The following protocol was used: GI identifiers of the homologous sequences identified by the BLAST search were used to retrieve UniProt IDs making use of a mapping file from iProClass database (<http://pir.georgetown.edu/pirwww/dbinfo/iproclass.shtml>). The iProClass is an integrated database for protein functional analysis, provides comprehensive descriptions of all proteins, with links to over 50 databases of proteins family, function, pathway, interaction, modification, structure, genome, ontology, literature and taxonomy.

RESULTS

Membrane proteins include integral and peripheral membrane proteins. The integral membrane proteins are a constitutive part of the membrane and can be classified into two categories: transmembrane proteins that span the entire membrane and integral monotopic proteins, which attach to the membrane from one side only. Peripheral membrane proteins are temporarily bound either to the lipid bilayer or to integral proteins by a combination of non-covalent interactions, such as hydrophobic and electrostatic interactions. *T. gondii* genome data is available at <http://Toxodb.org> (<http://Eupathdb.org>). In addition, a predicted proteome based on several gene predictions for *T. gondii* is available at EPIC-DB (<http://toro.aecom.yu.edu/biodefense/>). Theoretically, membrane proteins of *T. gondii* can be identified in these sequences using membrane segment prediction algorithms. However, the analysis is complicated by the fact that different protein predictions in *T. gondii* have about a 30% error rate in identifying coding regions (24, 31-33) and the predicted membrane proteins may be

expressed differently in different developmental stages. To experimentally validate gene predictions in *T. gondii* and comprehensively analyze the membrane proteome of *T. gondii*, we implemented three proteomics strategies as shown in Fig. 1. Strategy 1 and 2 target the whole membrane proteome, while strategy 3 targets a particular subset of the membrane proteome: the integral plasma membrane proteome.

One Dimensional SDS PAGE and NanoLC-MS/MS Analysis for *T. gondii* Membrane

Protein Extract

One dimensional (1D) SDS PAGE protein separation and in-gel tryptic digestion followed by reverse phase liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis (Strategy 1 in Fig. 1) is the most common and straightforward approach for the analysis of a complicated protein sample. The main advantage of 1D SDS PAGE separation over two-dimensional gel electrophoresis for membrane proteins is that many membrane proteins precipitate under isoelectric focusing conditions, but in general, most membrane proteins are soluble under SDS PAGE conditions (34, 35). After the membrane proteins were resolved on SDS PAGE, 40 to 45 gel bands were cut from each gel lane. Tryptic digestion was performed for each gel band after reduction and alkylation and the tryptic peptide mixture extracted from each band was subject to analysis by nanoLC-MS/MS. In the initial experiment, a wash step with a high salt and high pH buffer was not included in the membrane protein preparation procedure. A total of 1606 proteins were identified (see the green circle in Fig. 2A, $1169 + 437 = 1606$), of which 734 were identified as membrane proteins with one or more predicted transmembrane segments (see the green circle in Fig. 2B, $521 + 213 = 734$). These 734 proteins can be further grouped into 266 protein clusters (see the green circle in Fig. 2B, $178 + 88 = 266$; note that the number within a bracket presents the

number of protein clusters). The clustering approach groups together amino acid sequences even with very different lengths if the overlapping parts share more than 90% sequence identity (our clustering approach is described in detail in our recent paper, reference number 24). The protein clustering strategy was designed to group together alternative gene predictions of the same genomic regions, some of which may present possible splice variants and protein post-translational modifications, such as protease cleavages and covalent modifications.

High-salt buffer and alkaline buffer can disrupt non-covalent interactions between proteins which can then be used to wash away a large proportion of plasma membrane-associated cytosolic proteins (21, 36). Therefore, in the second experiment, we applied high salt and high pH buffer (1M KCl and 0.1 M Na₂CO₃) wash steps in the preparation of membrane proteins. Following extraction of the membrane fraction, the membrane protein extract was washed twice sequentially with 1M KCl and 0.1 M Na₂CO₃. In this experiment, 2938 proteins were identified (see the blue circle in Fig. 2A), of which 981 proteins (corresponding to 383 protein clusters) were predicted to have at least one transmembrane segment, indicating that this wash step had led to a remarkable increase in the number of membrane proteins identified (see the blue circle in Fig. 2B). There are 521 membrane proteins (in 178 protein clusters) that are common to both of the above experiments (Fig. 2B).

Relatively high abundance plasma membrane-associated cytosolic proteins, such as 2C-methyl-D-erythritol 2, 4-cyclodiphosphate synthase, cytochrome C oxidase, glycosyl transferase, were removed from the membrane protein fraction by washing with high salt and high pH buffer. The removal of abundant cytosolic proteins resulted in a decrease of protein dynamic range in the sample, permitting many more proteins to be identified by mass spectrometry, including H⁺-translocating inorganic pyrophosphatase TVP1, P-type ATPase, seven

transmembrane receptor domain-containing proteins and putative sodium hydrogen exchangers. H⁺-translocating pyrophosphatase (H⁺-PPases) has fourteen transmembrane segments and is the primary H⁺ pump that uses inorganic pyrophosphate (PPi) instead of ATP as an energy source. P-type ATPase and sodium hydrogen exchanger are predicted to have 11 and 13 transmembrane segments, respectively. Fig. 3A demonstrates the peptide coverage of a membrane protein (TgGlmHMM_2178), Ca²⁺ dependent ATPase with 11 predicted transmembrane segments. In total, 14 and 6 peptides (in red and blue colors) were identified with a Mascot peptide ion score higher than 60 or within a range of 49-60, respectively. Fig. 3B shows the MS/MS spectrum of a peptide (LLGCLDSLAADEK) of protein TgGlmHMM_2178. As shown in Fig. 3A, this peptide extends into the membrane. The zoom scan (inserted spectrum in Fig. 3B) clearly indicates that the peptide ion is doubly charged; and 83% of the *b* and *y* ions in the MS/MS spectrum match to the theoretical *b* and *y* ions, resulting in the unambiguous identification of this peptide.

TLSGE MudPIT Analysis for *T. gondii* Membrane Proteins

In Strategy 2 (Fig. 1), the three-layer “Sandwich” gel electrophoresis (TLSGE) method and Multi-dimensional chromatograph protein identification technology (MudPIT) were integrated for TLSGE MudPIT. Traditionally, MudPIT employs in-solution digestion of a complex protein mixture prior to two-dimensional liquid chromatography peptide separation. In TLSGE MudPIT a single in-gel digestion is needed before two-dimensional liquid chromatography. TLSGE MudPIT shares most of advantages of the conventional MudPIT alone, but has unique features well suited for membrane protein identification. Using this technique on the membrane protein preparation for *T. gondii*, using the same amount of organism as used for 1D gel LC-MS/MS,

without the high salt and high pH washes, 2938 proteins corresponding to 923 protein clusters were identified (see the green circle in Fig. 4A). Of these proteins, 1461 proteins were identified as membrane proteins with one or more predicted transmembrane segments, which grouped into 525 membrane protein clusters (green circle in Fig. 4B). When high salt and high pH wash steps were included in the sample preparation, 3116 proteins were identified (blue circle in Fig. 4A), of which 1428 were identified as membrane proteins (blue circle in Fig. 4B). These membrane proteins classified into 546 protein clusters (blue circle in Fig. 4B). There were 1825 proteins common to the two experiments (region of overlap in Fig. 4A), of which 893 proteins in 312 protein clusters are predicted with one or more transmembrane segments (region of overlap in Fig. 4B). In total, the two experiments led to identification of 4229 proteins from 1403 protein clusters, of which 1996 proteins from 759 protein clusters were found to be transmembrane proteins. In summary, the TLSGE MudPIT method identified 376 more transmembrane protein clusters than the 1D gel LC-MS/MS method.

Plasma Integral Membrane Proteome Analysis by Biotin-Directed Affinity Purification (BDAP) and NanoLC-MS/MS

To specifically target integral plasma membrane proteins located on the *T. gondii* cell surface, we implemented cell surface biotinylation and affinity purification (20, 21, 37, and 38), using the same amount of organism as used for 1D gel LC-MS/MS (see strategy 3 in Fig.1). In total, 1029 proteins in 304 protein clusters were identified (the green circle in Fig. 5 A), of which 329 proteins have predicted transmembrane segments and grouped into 118 membrane protein clusters (the green circle in Fig. 5B). Besides many hypothetical membrane proteins, these include many

known or putative plasma integral membrane proteins in *T. gondii* with one or multiple predicted transmembrane segments, such as rhoptry proteins 4, 5, 13, 14 and 18; rhoptry neck proteins 1, 2 and 3; dense granule protein GRA3, 6, and 7; ADP/ATP carrier; calcium ATPase SERCA-like protein; Rab11; Ca²⁺ ATPase; facilitative glucose transporter GT1; dihydrolipoamide acetyltransferase; ATP-binding cassette protein subfamily B member 3; H⁺ translocating inorganic pyrophosphatase TVP1 protein; oxoglutarase/ malate translocator protein; vacuolar H⁺ translocating ATPase subunit A; vacuolar ATP synthase catalytic subunit A; oxalate formate antiporter; and NMDA receptor glutamate-binding chain.

Comparison of the Three Proteomics Strategies

Fig. 5 summarizes all the identified proteins and transmembrane proteins by the three approaches. In total, 2241 *T. gondii* membrane proteins from 841 protein clusters were identified (Fig. 5B). Approximately 66% were identified by at least two peptide matches, each with a Mascot score of 49 or higher. Meanwhile 15.4 % were identified by a single peptide match with a score larger than 60, and 18.8% by a single peptide match with a score between 49 and 60. Fig. 6A shows the distribution of peptide Mascot scores for all the identified transmembrane proteins by all the three methods. The majority (96%) of these peptides had a Mascot score greater than 60 and were repeatedly found in three of the different proteomics experiments. The distribution of identified membrane proteins across the number of predicted transmembrane segments is shown in Fig. 6B. Approximately 62 % of all identified transmembrane proteins have two or more predicted transmembrane segments and 20.8 % of all identified transmembrane proteins have 7 or more transmembrane segments. One protein (52.m01648), which has 22 predicted transmembrane

segments, was identified as the adenylate and guanylate cyclase catalytic domain-containing protein (Supplementary Table 1).

We investigated the overlap of the identified proteins as well as membrane proteins among the three approaches (Fig.5). The overlap between 1D gel LC-MS/MS and TLSGE MudPIT reveals that 959 membrane proteins from 358 protein clusters were common to both methods (Fig. 5B). These common membrane proteins represent 48% and 80.3% of the membrane proteins identified by TLSGE MudPIT or 1D gel LC-MS/MS, respectively. 285 identified membrane proteins from 98 protein clusters were common to all three methods, which were 23.9%, 14.3% and 86.6 % of the membrane proteins identified by 1D gel LC-MS/MS, TLSGE MudPIT and BDAP LC-MS/MS approaches, respectively. The two former methods target the whole membrane proteome, whereas BDAP LC-MS/MS targets a particular fraction of the membrane proteome: integral plasma membrane proteins. Only 10 membrane proteins from 4 protein clusters were exclusively identified by BDAP LC-MS/MS. The combination of 1D gel LC-MS/MS and TLSGE MudPIT analysis was able to identify about 97% of integral plasma membrane proteins. However, BDAP LC-MS/MS analysis is important, as it allows screening specifically for plasma integral membrane proteins, many of which are potential drug targets. Supplementary Table 2 A to G list all the identified proteins in regions of “a” to “g” in Fig. 5A. Supplementary Table 3 A to G list all the identified transmembrane proteins in regions of “a” to “g” in Fig.5B.

Analysis of the Proteins Identified by the Combined Strategies

Supplementary Table 1 contains all of the identified *T. gondii* membrane proteins with

protein accession key, number of predicted transmembrane segments, number of peptides assigned to the protein, the experiments in which the protein was found, as well as name or function of the protein. By single clicking the web linked sequence keys in supplementary Table 1, one can retrieve the detailed information of each identified transmembrane protein cluster from the EPIC-DB database (<http://toro.aecom.yu.edu/cgi-bin/biodefense/main.cgi>).

All proteins identified with one or more predicted transmembrane segments were subjected to Gene Ontology (GO) analysis (39). As demonstrated in Fig. 7A, 23% of the proteins are classified as membrane proteins, 21% are integral to membrane, 3% are plasma membrane proteins, as well as 3% as endoplasmic reticulum membrane proteins. Biological process GO analysis indicates that a large number of the identified membrane proteins (15%) are involved in transport, including 3% and 2% in cation and ion transport, respectively (Fig. 7B). Furthermore, many of the identified membrane proteins (4%) are involved in ATP biosynthetic processes and 7% in metabolic processes. Approximately 4% of the identified membrane proteins function in proteolytic and protein phosphorylation processing (Fig. 7B). In molecular function analysis, many identified membrane proteins are characterized as ATP, nucleotide, protein or metal ion binding molecules (Fig. 7C). Many are predicted to have a membrane-related enzymatic activity, such as ATPase activity coupled to transmembrane movement of ions or other substances or phosphorylative mechanism, as well as hydrolase, transferase and oxidoreductase activities. GPI modification site prediction analysis reveals 29 proteins highly likely to be glycosylphosphatidylinositol (GPI) anchored proteins, such as surface antigen 43, a SRS domain containing protein and GPI-anchored surface BSR4-related antigen. In addition, many proteins are likely to be GPI anchored, for instance, SAG2-related antigen SAG2E, signal recognition particle 54Kda protein, sporozoite-specific SAG protein and so on. Interestingly, many of these

predicted GPI anchored proteins are hypothetical proteins.

It was striking that more than 42% of membrane proteins identified in the current study are hypothetical proteins. Furthermore, half of these hypothetical proteins appear to be unique to *T. gondii*. For example, a hypothetical membrane protein (TgGlmHMM_0264) having eight predicted transmembrane segments, was identified by all three proteomics approaches with 8 peptide matches, four with Mascot peptide ion scores greater than 60 and four with Mascot ion scores between 49 and 60. The negative outcome of BLAST searches and GO annotations suggest that this protein is of unknown function and unique to *T. gondii*.

Since *T. gondii* parasites were cultured in human foreskin fibroblasts, there is a possibility that the membrane protein preparation of *T. gondii* may contain human protein contaminants. To remove the human protein hits, we performed Mascot searches for all MS/MS data against a protein sequence database combining the human proteome and the predicted *T. gondii* protein sequence data sets. In total, 1161 proteins in 662 clusters were identified as human proteins. 1045 out of these 1161 human proteins do not have any peptide hits that match *T. gondii* proteins, which suggests that they are unique to human. The rest of the 116 human proteins have one or more peptide hits in common with *T. gondii* proteins, however, only two (NCBI gi number 115511024 and 153945715) have predicted transmembrane segments.

All the identified *T. gondii* proteins with at least one predicted transmembrane segment are listed in Supplementary Table1. They represent approximately 25.2 % of the predicted membrane proteome (8897 predicted membrane proteins which group into 3858 clusters) from the *T. gondii* genome. Since in this study we investigated only the tachyzoite life stage, the coverage of the *T. gondii* membrane proteome by these experiments is quite high. All the data for *T. gondii* membrane proteome analysis as well as bioinformatics analysis have been deposited on the

Experimental Proteomics DataBase for *Toxoplasma gondii* in the Einstein Biodefense Proteomics Research Center (<http://toro.aecom.yu.edu/cgi-bin/biodefense/main.cgi>) and the data provided to ToxodB (<http://ToxodB.org>) which is part of EuPathdB.

DISCUSSION

Membrane proteins are critical players in various physiological processes, such as regulation of intracellular pH; the control of cell growth and proliferation; transepithelial absorption and secretion of sodium, hydrogen, bicarbonate, chloride and organic anions; and the metabolic response to hormones such as insulin and glucocorticoids (40). Many diseases or disorders are related to the functions of membrane proteins. A large portion of current pharmaceutical drug targets are membrane proteins. Similar to other organisms, about 30% of proteins encoded in the genome of *T. gondii* are predicted to be membrane proteins. However, a great number of them are unknown or have not been characterized although the genome sequencing of this parasite has been completed for several years. Expression profiling of *T. gondii* membrane proteins at different developmental stages and accurate membrane gene annotation is important for understanding the invasive mechanisms and the interaction between *T. gondii* and their hosts. We report here that by utilizing three proteomic strategies, 1D gel LC-MS/MS, TLSGE MudPIT and BDAP LC-MSMS, we were able to identify and validate more than 2000 membrane proteins of *T. gondii* tachyzoites, classified into 841 protein clusters with a requirement of 90% sequence identity threshold in each cluster. Since the tandem mass spectra data were searched against the hypothetical *T. gondii* proteome generated through a combination of computationally predicted proteins from TigrScan, TwinScan, GlimmerHMM, Release 4.3

(ToxodB) and ME49 (Release 6.0 ToxodB), and the available experimental *T. gondii* sequences from the NCBI non-redundant protein database, a protein clustering strategy is essential to remove redundancy. In addition, the protein sequence similarity clustering strategy also provides insights into potential splicing products within the genome. Each non-redundant cluster may represent a single sequence or multiple sequences. It was found that the non-redundant protein clusters generated for *T. gondii* have an average size of 2.02 sequences (24). Multiple sequences in a cluster may be truly redundant sequences or are the shorter forms generated from the longer form due to naturally occurring enzymatic process.

Currently, MS/MS database searching is the primary approach for protein identification in large scale proteomic analysis. False positive matches are known to occur (41). Randomized decoy database search strategies are frequently used to estimate the False Discovery Rate (FDR) of peptide identification (42, 43). We implemented the Mascot search tool with an automatic decoy search feature to perform MS/MS database searches. In this study, the Mascot peptide ion score threshold for accepting a MS/MS match to a certain peptide was set at 49 (individual peptide ion scores > 49 indicate identity or extensive homology, $p < 0.05$). This threshold led the overall FDR of peptide identification to be 1.1 %. For the proteins identified with a single peptide match, additional manual checks were performed, to see if the adjusted observed peptide mass after removing the systematic error of mass spectrometry analysis is within the range of the theoretical mass plus or minus 100 ppm (this mass tolerance falls in the specific internal mass error of LTQ ion trap mass spectrometer), if the fragment ion mass tolerance is within 0.5 Da, if the MS/MS fragment ions have both *b* and *y* ions, and if 80% of the most intensity fragment ions match to either *b* or *y* ions. A recently published work investigated the effect of the “two-peptide” rule on protein identification [44]. The authors demonstrated that the “two-peptide” rule decreases the

number of protein identifications in the target database more significantly than in the decoy database, resulting in increased FDR, when compared to the case when proteins identified by only a single peptide match are not discarded (44). A large fraction of proteins may be missed by the “two-peptide” rule in proteomic experiments (44). In our case, about 34% of the 841 membrane protein clusters were identified by only a single peptide match. Most of these single peptide matches were repeatedly found in the experiments carried out independently by the three different membrane proteomic approaches, further supporting that these single peptide matched proteins were present in the organism. Therefore, we included those membrane proteins identified with one peptide match (Supplementary Table 1). The single-peptide-hit identified membrane proteins are most likely to be those proteins, which are very low abundance. In addition, integral membrane proteins have a significant percentage of sequence embedded in the membrane and this may also account for the single peptide matches observed. The commonly utilized chromatography of LC-MS/MS for proteomic study is generally suited to the soluble peptide segments of integral membrane proteins and cannot efficiently access membrane-embedded peptides (45). Therefore, the identification sequence coverage of an integral membrane protein is often lower than that of a cytosolic protein. Because of these issues and the fact that different protein predictions in *T. gondii* have been shown to each have about a 30% error rate in identifying coding regions (24), it is important to have proteomics strategies in place that focus on recognizing these low abundant proteins and identifying the membrane-embedded peptides.

To achieve a global analysis of the membrane proteome of even such a small organism, one cannot rely on one approach alone. Because the proteome is much more complicated and less defined than the genome, more comprehensive proteomic strategies are essential. We applied three approaches, orthogonal to each other, in order to decipher the membrane proteome of *T.*

gondii. The BDAP LC-MS/MS method specifically targets integral plasma membrane proteins; while 1D gel LC-MS/MS and TLSGE MudPIT methods targeted the whole membrane proteome. The common membrane proteins identified by using all three approaches represent 13% of all the identified membrane proteins but 87% of membrane proteins identified by BDAP LC/MSMS method (Fig. 7). The membrane proteins identified either by 1D gel LC-MS/MS or TLSGE MudPIT share about 43% of all the identified membrane proteins from *T. gondii* tachyzoites. Both approaches uniquely identified a large number of membrane proteins, and the unique membrane proteins identified by TLSGE MudPIT was about 3 times more than those identified by 1D gel LC-MS/MS. These data demonstrate that none of these approaches can completely replace the other, which is especially true for the 1D gel LC-MS/MS and TLSGE MudPIT approaches. This also suggests complementary proteomic strategies must be employed in different developmental stages of an organism to achieve more complete proteomic coverage.

High salt and high pH buffer (sodium carbonate) can be used to strip membrane-associated proteins and enrich membrane proteins (46-52). *T. gondii* has unique structures associated with the cell membrane including the pellicle, apical rings, polar rings, conoid, rhoptries, and micronemes, as compared to other cells. Its outer covering pellicle consists of plasmalemma and two closely applied membranes that form an inner membrane complex (IMC). Between the parasite plasma membrane and IMC, there are the myosins A motor complex and the micronemal protein-host receptor complex (via aldolse/F-actin) at the parasite pellicule (53). Furthermore, beneath the IMC, 22 subpellicular microtubules run longitudinally along all length of the cell. The complicated and tight interactions among these structures often lead to the contamination of cytoskeleton or cytoplasm proteins in membrane preparation. Washing with high salt and high pH buffer substantially removed the abundant membrane-associated proteins from our *T. gondii*

membrane prep thus, making it possible to additionally identify a large number of membrane proteins. Interestingly, high salt and high pH buffer wash appeared to impact the identification of membrane proteins in 1D gel LC-MS/MS more so than in the TLSGE MudPIT strategy. When combining the proteins identified by experiments with or without high salt and high pH buffer washes, the latter approach identified 288 more membrane protein clusters than the former. However, with such washes, 1D gel LC-MS/MS identified 40% more membrane protein clusters than without these washes, whereas, for TLSGE MudPIT experiment, the number of identified membrane protein clusters with these wash steps is comparable to that without these washes (cf. Fig. 2 and Fig. 4). This phenomenon probably indicates that the dynamic range of 1D gel LC-MS/MS analysis is smaller than that of TLSGE MudPIT analysis. However, although a high salt and high-pH buffer wash is able to significantly decrease the abundance of membrane-associated proteins, this harsh and stringent wash may also decrease the abundance of some membrane proteins. This has been observed previously by other researchers (54) and is also strongly supported by our proteomic data from both 1D gel LC-MS/MS and TLSGE MudPIT experiments. Peripheral membrane proteins dissociate from transmembrane proteins following a treatment with a solution having an elevated pH or high salt concentrations. Also, the high salt and high pH conditions might denature some membrane proteins, possibly releasing some of them into the wash solution.

To our knowledge, this work represents the most comprehensive analysis of the membrane proteome of a microorganism by integrating three proteomic approaches with different sample preparation methods and computational bioinformatics investigations. This integrative analysis identified thousands of membrane proteins in *T. gondii*. Many are known membrane proteins, including proteins from plasma membrane and rhoptry, microneme, dense granule,

parasitophorous vacuole organelles. Many membrane proteins putatively function as membrane enzymes, as transporters or carriers, ion exchangers, surface antigens, receptors, structural membrane-anchoring proteins, as well as chaperone proteins (e.g., disulfide isomerases and regulators of vesicular traffic like ADP ribosylation factors). The most intriguing finding is that a large number of transmembrane proteins identified in this work are hypothetical proteins. Furthermore, approximately 50% of these hypothetical proteins have no GO annotations, suggesting that these sequences are most likely unique to *T. gondii* parasites or the Apicomplexa. These unique and novel membrane proteins may represent potential drug targets and further systematic analysis of the membrane proteome of *T. gondii* will provide additional fundamental insights into the biology and molecular mechanisms underlying the pathophysiology of infection of this obligate intracellular parasite.

ACKNOWLEDGMENTS

This article is dedicated to the memory of Professor George Orr, Ph.D., deceased November 2005. Dr. George Orr was the founder of the Einstein Biodefense Proteomics Research Center. This work was supported by NIH/NIAID Contract HHSN266200400054C and NIH AI39454 (LMW).

REFERENCES

1. Ajioka, J.W., Fitzpatrick, J.M., and Reitter CP. (2001) *Toxoplasma gondii* genomics: shedding light on pathogenesis and chemotherapy. *Expert Rev. Mol. Med.* **3**, 1-19

2. Lang, C., Groß, U., and Lüder, C.G.K. (2007) Subversion of innate and adaptive immune responses by *Toxoplasma Gondii*. *Parasitol. Res.* **100**, 191-203
3. Dubey, J.P. Toxoplasmosis- a waterborne zoonosis. (2004) *Vet. Parasitol.* **126**, 57-72
4. Dawson, D. Foodborne protozoan parasites. (2005) *Int. J. Food Microbiol.* **103**, 207-227
5. Carey, K.L., Westwood, N.J., Mitchison, T.J., and Ward, G.E. (2004) A small-molecule approach to studying invasive mechanisms of *Toxoplasma gondii*. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 7433-7438
6. Sibley, L.D., and Ajioka, J.W. (2008) Population Structure of *Toxoplasma gondii*: Clonal Expansion Driven by Infrequent Recombination and Selective Sweeps. *Annu. Rev. Microbiol.* **62**, 329-351
7. Correa, D., Canedo-solares, I., Ortiz-alegeria, L.B., Caballero-ortega, H., and Rico-torres, C.P. (2007) Congenital and acquired toxoplasmosis: diversity and role of antibodies in different compartments of the host. *Parasite Immunol.* **29**, 651-660
8. Kasper, L., Courret, N., Darche, S., Luangsay, S., Mennechet, F., Minns, L., Rachinel, N., Ronet, C., and Buzoni-Gatel, D. (2004) *Toxoplasma gondii* and mucosal immunity. *Int. J. Parasitol.* **34**, 401-409
9. Kravetz, J.D., and Federman, D.G. (2005) Toxoplasmosis in pregnancy. *Am. J. Med.* **118**, 212-216
10. Boothroyd, J.C., and Dubremetz, J. (2008) Kiss and spit: the dual roles of *Toxoplasma* rthroptries. *Nat. Rev. Microbiol.* **6**, 79-88
11. Carruthers, V., and Boothroyd, J.C. (2007) Pulling together: an integrated model of *Toxoplasma* cell invasion. *Curr. Opin. Microbiol.* **10**, 83-89.
12. Lingelbach, K., and Joiner, K.A. (1998) The parasitophorous vacuole membrane surrounding

plasmodium and Toxoplasma: an unusual compartment in infected cells. *J. Cell Sci.* **111**, 1467-1475

13. Mital, J., Meissner, M., Soldati, D., Ward, G.E. (2005) Conditional expression of Toxoplasma gondii apical membrane antigen-1(TgAMA1) demonstrates that TgAMA1 plays a critical role in host cell invasion. *Mol. Biol. Cell* **16**, 4341-4349

14. Bouchot, A., Millot, J.M., Charpentier, S., Bonhomme, A., Villena, I., Aubert, D., and Pinon, J.M. (2001) Membrane potential changes after infection of monocytes by Toxoplasma gondii. *Int. J. Parasitol.* **31**, 1114-1120

15. Lekutis, C., Ferguson, D.J.P., Grigg, M.E., Camps, M., and Boothroyd JC. (2001) Surface antigens of Toxoplasma gondii: variations on a theme. *Int. J. Parasitol.* **31**, 1285-1292

16. Meissner, M., Reiss, M., Viebig, N., Carruthers, V.B., Toursel, C., Tomavo, S., Ajioka, J.W., and Soldati, D. (2002) A family of transmembrane microneme proteins of *Toxoplasma gondii* contain EGF-like segments and function as escorts. *J. Cell Sci.* **115**, 563-574

17. Loo, R.R.O., Dales, N., and Andrews, P.C. (1994) Surfactant effects on protein structure examined by electrospray ionization mass spectrometry. *Protein Sci.* **3**, 1975-1983

18. Galeva, N., and Altermann, M. (2002) Comparison of one-dimensional and two-dimensional gel electrophoresis as a separation tool for proteomic analysis of rat liver microsomes: Cytochromes P450 and other membrane proteins. *Proteomics* **2**, 713-722

19. Ferro, M., Salvi, D., Riviere-Rolland, H., Vermat, T., Seigneurin-Berny, D., Grunwald, D., Garin, J., Joyard, J., and Rolland, N. (2002) Integral membrane proteins of the chloroplast envelope: identification and subcellular localization of new transporters. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 1148-11492

20. Zhao, Y., Zhang, W., Kho, Y., and Zhao, Y. (2003) Capillary HPLC/MS spectrometric

analysis of proteins from affinity-purified plasma membrane. *Anal. Chem.* **75**, 3751-3757

21. Zhao, Y., Zhang, W., Kho, Y, and Zhao, Y. (2004) Proteomic analysis of integral plasma membrane proteins. *Anal. Chem.* **76**, 1817-1823

22. Lu, X., and Zhu, H. (2005) Tube-Gel Digestion. *Mol. Cell. Proteomics* **4**, 1948-1958

23. Liu, T., Martin, A.M., Sinai, A.P., and Lynn, B.C. (2008) Three-layer Sandwich gel electrophoresis: A method of salt removal and protein concentration in proteome analysis. *J. Proteome Res.* **7**, 4256-4265

24. Dybas, J.M., Madrid-Aliste, C.J., Che, F.Y., Nieves, E., Rykunov, D., Angeletti, R.H., Weiss, L.M., Kim, K., and Fiser, A. (2008) Computational analysis and experimental validation of gene predictions in *Toxoplasma gondii*. *PLoS One* **3**, e3899

25. Madrid-Aliste, C.J., Dybas, J.M., Angeletti, R.H., Weiss, L.M., Kim, K., Simon, I., and Fiser, A. (2009) EPIC-DB: a proteomics database for studying Apicomplexan organisms. *BMC Genomics* **10**, 38

26. Li, W.Z., Jaroszewski, L., and Godzik, A. (2001) Clustering of highly homologous sequences to reduce the size of large protein database. *Bioinformatics* **17**, 282-283

27. Li, W.Z., Jaroszewski, L., and Godzik, A. (2002) Tolerating some redundancy significantly speeds up clustering of large protein databases. *Bioinformatics* **18**, 77-82

28. Li, W.Z., and Godzik A. (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**, 1658-1659

29. Käll, L., Krogh, A., and Sonnhammer, E.L.L. (2004) A combined transmembrane topology and signal peptide prediction method. *J. Mol. Biol.* **338**, 1027-1036

30. Pierleoni, A., Martelli, P.L., and Casadio, R. (2008) PredGPI: a GPI-anchor predictor. *BMC Bioinformatics* **9**, 392

31. Lu, F., Jiang, H., Ding, J., Mu, J., Valenzuela, J.G., Ribeiro, J. M.C., and Su, X. (2007) cDNA sequences reveal considerable gene prediction inaccuracy in the *Plasmodium falciparum* genome. *BMC Genomics* **8**, 255
32. Guigo, R., Flicek, P., Abril, J.F., Reymond, A., Lagarde, J., Denoeud, F., Antonarakis, S., Ashburner, M., Bajic, V.B., Birney, E., Castelo, R., Eyraes, E., Ucla, C., Gingeras, T.R., Harrow, J., Hubbard, T., Lewis, S.E., and Reese, M.G. (2006) EGASP: the human ENCODE Genome Annotation Assessment Project. *Genome Biol.* **7**, (Suppl 1) S2.1-31
33. Nielsen, P., and Krogh, A. (2005) Large-scale prokaryotic gene prediction and comparison to genome annotation. *Bioinformatics* **21**, 4322-4329
34. Wu, C.C., Yates III, J.R. (2003) The application of mass spectrometry to membrane proteomics. *Nat. Biotechnol.* **21**, 262-267
35. Xiong, Y., Chalmers, M.J., Gao, F.P., Cross, T.A., and Marshall, A.G. (2005) Identification of *mycobacterium tuberculosis* H37Rv integral membrane proteins by one-dimensional gel electrophoresis and liquid chromatography electrospray ionization tandem mass spectrometry. *J. Proteome Res.* **4**, 855-861
36. Speers, A.E., and Wu, C.C. (2007) Proteomics of integral membrane proteins-Theory and Application. *Chem. Rev.* **107**, 3687-3714
37. Nunomura, K., Nagano, K., Itagaki, C., Taoka, M., Okamura, N., Yamauchi, Y., Sugano, S., Takahashi, N., Izumi, T., and Isobe, T. (2005) Cell Surface Labeling and Mass Spectrometry Reveal Diversity of Cell Surface Markers and Signaling Molecules Expressed in Undifferentiated Mouse Embryonic Stem Cells. *Mol. Cell. Proteomics* **4**, 1968–1976
38. Sostaric, E., Georgiou, A.S., Wong, C.H., Watson, P.F., Holt, W.V., and Fazeli, A. (2006) Global Profiling of Surface Plasma Membrane Proteome of Oviductal Epithelial Cells. *J.*

Proteome Res. **5**, 3029–3037

39. Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K. S., Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P/, Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., and Sherlock, G. (2000) Gene Ontology: tool for the unification of biology. *Nat. Genetics* **25**, 25 – 29
40. Mahnensmith, R.L., and Aronson, P.S. (1985) The plasma membrane sodium-hydrogen exchanger and its role in physiological and pathophysiological processes. *Circ. Res.* **56**, 773-788
41. Cargile, B. J., Bundy, J.L., and Stephenson, J.L.Jr. (2004) Potential for false positive identifications from large databases through tandem mass spectrometry. *J. Proteome Res.* **3**, 1082-1085
42. Elias, J.E., and Gygi, S.P. (2007) Target-decoy search strategy for increased confidence in large-scale protein identifications by mass spectrometry. *Nat. Methods* **4**, 207-214
43. Kall, L., Storey, J.D., MacCoss, M.J., and Noble, S.W. (2008) Assigning significance to peptides identified by tandem mass spectrometry using decoy databases. *J. Proteome Res.* **7**, 29-34
44. Gupta, N., and Pevzner, P.A. (2009) False discovery rates of protein identifications: a strike against the two-peptide rule. *J. Proteome Res.* **8**, 4173-4181
45. Speers, A.E., Blackler, AR., and Wu, C.C. (2007) Shotgun analysis of integral membrane proteins facilitated by elevated temperature. *Anal. Chem.* **79**, 4613-4620
46. Pedersen, S.K., Harry, J.L., Sebastian, L., Baker, J., Traini, M.D., McCarthy, J.T., Manoharan, A., Wilkins, M.R., Gooley, A.A., Righetti, P.G., Packer, N.H., Williams, K.L., and Herbert, B.R. (2003) Unseen Proteome: Mining Below the Tip of the Iceberg To Find Low Abundance and Membrane Proteins. *J. Proteome Res.* **2**, 303-311

47. Zhang, L.J, Wang, X., Peng, X., Wei, Y.J., Cao, R., Liu, Z., Xiong, J.X., Yin, X., Ping, C., and Liang, S. (2006) Proteomic analysis of low-abundant integral plasma membrane proteins based on gels. *Cell Mol. Life Sci.* **63**, 1790-1804
48. Moebius, J., Zahedi, R.P., Lewandrowski, U., Berger, C., Walter, U., and Sickmann, A. (2005) The Human Platelet Membrane Proteome Reveals Several New Potential Membrane Proteins. *Mol. Cell. Proteomics* **4**, 1754-1761
49. Schirmer, E. C., Florens, L., Guan, T., Yates III, J.R., and Gerace L. (2003) Nuclear membrane proteins with potential disease links found by subtractive proteomics. *Science* **301**, 1380-1382
50. Fujiki, Y., Hubbard, A.L., Fowler, S., and Lazarow, P.B. (1982) Isolation of intracellular membranes by means of sodium carbonate treatment :application to endoplasm is reticulum. *J. Cell Biol.* **93**, 97-102
51. Mattow, J., Siejak, F., Hagens, K., Schmidt, F., Koehler, C., Treumann, A., Schaible, U.E., and Kaufmann, S.H.E. (2007) An improved strategy for selective and efficient enrichment of integral plasma membrane proteins of mycobacteria. *Proteomics* **7**, 1687-1701
52. Ruth, M. C., Old, W. M., Emrick, M. A., Meyer-Arendt, K., Aveline-Wolf, L. D., Pierce, K. G., Mendoza, A. M., Sevinsky, J. R., Hamady, M., Knight, R. D., Resing, K. A., and Ahn, N. G. (2006) Analysis of membrane proteins from human chronic myelogenous leukemia cells: comparison of extraction methods for multidimensional LC-MS/MS. *J. Proteome Res.* **5**, 709-719
53. Soldati, D., Meissner, M. (2004) Toxoplasma as a novel system for motility. *Curr. opin. cell boil.* **16**, 32-40
54. Mastrogiacomo, A., Kohan, S.A., Whitelegge, J.P., Gundersen, C.B. (1998) Intrinsic membrane association of Drosophila cysteine string proteins. *FEBS Lett.* **436**: 85-91

FIGURE LEGENDS

Figure 1. Scheme describing experimental strategies for the comprehensive analysis of the membrane proteome of *T. gondii*. Strategy 1 involved in-gel tryptic digestion from bands cut from a 1D SDS gel. Strategy 2 utilized the three-layer sandwich electrophoresis method (23) where proteins were concentrated in the middle, 10-12% polyacrylamide (PA) layer, followed by in-gel digestion and strong cation exchange (SCX) fractionation. In strategy 3 cell surface proteins were biotin labeled and affinity purified. NanoLC-MS/MS and Mascot searches were performed after all 3 strategies.

Figure 2. Venn diagrams showing the number of proteins identified by Mascot having peptide ion scores of 49 or greater from the 1 D gel LC-MS/MS experiment (Strategy 1). A, Total number of proteins identified with or without Na₂CO₃ and KCl washes. B, Number of membrane proteins identified in A having one or more predicted transmembrane segments with or without Na₂CO₃ and KCl washes. The numbers in the square brackets refer to the number of protein clusters. The expression of “w/” means “with” and “w/o” means “without”. Proteins are listed in Supplementary Tables

Figure 3. (A) Amino acid sequence diagram of the transmembrane protein Ca²⁺ dependent ATPase (TgGlmHMM_2178). Peptides in red were identified by LC-MS/MS with a Mascot score of 60 or more. Peptides in blue are identified by LC-MS/MS with a Mascot score in a range of 49-60. The transmembrane segments were predicted with Phobius program. **(B) MS/MS spectrum of the tryptic peptide LLGCLDSLAADEK,** transmembrane region 2 in red, identified from all the three proteomic strategies. This peptide ion at *m/z* 702.84 (monoisotopic) is

doubly charged (see inserted zoom scan). The observed peptide parent mass is 1403.66 Da and the theoretical peptide monoisotopic mass is 1403.70. The major ions in this MS/MS spectrum comprise 83% of the predicted peptide's b and y ions. Note that the C residue is carboxamidomethyl cysteine.

Figure 4. Venn diagrams showing the number of proteins identified by Mascot having peptide ion scores of 49 or greater from the TLSGE MudPIT experiment (Strategy 2). A, Total number of proteins identified with or without Na₂CO₃ and KCl washes. B, Number of membrane proteins identified in A having one or more predicted transmembrane segments with or without Na₂CO₃ and KCl washes. The numbers in the square brackets refer to the number of protein clusters. The expression of “w/” means “with” and “w/o” means “without”. Proteins are listed in Supplementary Tables.

Figure 5. Venn diagrams of proteins identified from all 3 strategies, 1D gel LC-MS/MS, biotin labeling and TLSGE MudPIT. A, Total number of proteins identified with or without Na₂CO₃ and KCl washes. Number within square bracket refers to the number of protein clusters. Proteins in each region of “a” to “g” are listed in Supplementary Table 2A to G; B, Numbers of identified membrane proteins from A having one or more predicted transmembrane segments with or without Na₂CO₃ and KCl washes. Number within square bracket refers to the number of protein clusters. Proteins in each region of “a” to “g” are listed in Supplementary Table 3 A to G.

Figure 6. (A), Distribution of the number of identified peptides from Mascot searches vs Mascot peptide ion score. (B), Distribution of the number of identified proteins vs the

number of transmembrane segments. Results were obtained from all 3 strategies (Fig. 1).

Individual peptide ion scores = or > 49 indicate identity or extensive homology, $p < 0.05$. For the first bar, Mascot ion score region is 49 to 60. The width in x axis of rest of bar is 10, for example, for the second bar, the Mascot ion score region is 60 to 70. Proteins are listed in Supplementary Tables.

Figure 7. Pie charts of functional distribution per cellular component (A), biological processes (B) and (C) molecular function from the membrane proteins identified by all three proteomic strategies (Fig. 1). Numbers represent the percentage from the total of identified membrane proteins. Proteins are listed in Supplementary Tables.




Fig.1





Fig.2






Fig.3B




Fig.4




Fig.5





Fig.6




Fig.7